arXiv:0705.4080v1 [math.DS] 28 May 2007

Aperiodic substitutional systems and their Bratteli diagrams

S. Bezuglyi
Institute for Low Temperature Physics, Kharkov, Ukraine
bezuglyi@ilt.kharkov.ua

J. Kwiatkowski
Nicolaus Copernicus University, Toruń, Poland
jkwiat@mat.uni.torun.pl

K. Medynets*
Institute for Low Temperature Physics, Kharkov, Ukraine
medynets@ilt.kharkov.ua

Abstract

In the paper we study aperiodic substitutional dynamical systems arisen from non-primitive substitutions. We prove that the Vershik homeomorphism φ of a stationary ordered Bratteli diagram is homeomorphic to an aperiodic substitutional system if and only if no restriction of φ to a minimal component is homeomorphic to an odometer. We also show that every aperiodic substitutional system generated by a substitution with nesting property is homeomorphic to the Vershik map of a stationary ordered Bratteli diagram. It is proved that every aperiodic substitutional system is recognizable. The classes of m-primitive substitutions and associated to them derivative substitutions are studied. We discuss also the notion of expansiveness for Cantor dynamical systems of finite rank.

1 Introduction

During last decade, minimal homeomorphisms of a Cantor set (Cantor minimal systems, in other words) have been thoroughly studied in many papers. The most powerful tool in the study of Cantor minimal systems is the concept of Bratteli diagrams. It was shown in [HPS] that every minimal homeomorphism is topologically conjugate to the Vershik map of an ordered simple Bratteli diagram. Such a realization of minimal homeomorphisms allowed one to prove many deep results clarifying properties and orbit structure of minimal Cantor

---

*The third named author was supported by INTAS YSF-05-109-5315.

systems. We mention here the study of orbit equivalence and affability of Cantor minimal systems which was conducted in papers by Glasner, Giordano, Matui, Putnam, Skau, Weiss, and others (see [GPS1], [GPS2], [GPS3], [GMPS], [GW]). Another meaningful usage of Bratteli diagrams was given in the papers [For] and [DHS]. They answered the natural question of the description of the class of minimal homeomorphisms which can be represented by Bratteli diagrams of the simplest form, i.e., by stationary Bratteli diagrams. It turns out that this class is constituted by minimal substitutional systems and odometers.

Motivated by these remarkable achievements, we are interested in the following question: Is the assumption of minimality crucial in proving these results? In the paper [BDM], we considered aperiodic Cantor systems and proved the existence of Kakutani-Rokhlin partitions for them. In contrast to the minimal case, we cannot start with an arbitrary clopen set to produce a Kakutani-Rokhlin partition. Nevertheless, it was proved in [Med] that, given an aperiodic homeomorphism T, there exists a sequence of nested Kakutani-Rokhlin partitions which generate the topology. Thereby we showed that every aperiodic homeomorphism can be realized as the Vershik map of an ordered Bratteli diagram (see details in Section 2). However, the structure of Bratteli diagrams for aperiodic homeomorphisms is still unclear — in comparison with the minimal case where each simple Bratteli diagram can be given an order that defines a continuous Vershik map [HPS], not every Bratteli diagram admits, in general, such an ordering [Med].

The primary goal of this work is the study of aperiodic (non-minimal) substitutional systems and finding explicit descriptions of their Bratteli-Vershik models. To the best our knowledge, non-primitive substitutions have not been systematically studied yet. In our study we are mostly based on two articles. The first one is the paper by Durand, Host, and Skau [DHS] where the Bratteli diagrams for primitive substitutional dynamical systems were thoroughly studied. The second one is the very recent work by Downarowicz and Maass [DM] where the authors suggested a very fruitful idea of coding of dynamics by means of the so-called j-symbols. In other words, this approach gives a symbolic interpretation of the technique of Kakutani-Rokhlin partitions and Bratteli diagrams. The basic references to the study of substitutional dynamical systems are [Fog] and [Que]. We also refer the reader to the book [Kur] for a comprehensive exposition of the symbolic dynamics.

We will use the standard notation of the theory of substitutional systems. Denote by A a finite alphabet and by A^+ the set of all non-empty words over A. Let $\sigma : A \to A^+$ be a substitution. We denote by A_l the set of all letters $a \in A$ such that $|\sigma^n(a)| \to \infty$ as $n \to \infty$. Let $A_s = A \setminus A_l$. We say that a substitution σ has a *nesting property* if at least one of the following two conditions holds: (1) for every $a \in A_l$ the word $\sigma(a)$ starts with a letter from A_l; (2) for every $a \in A_l$ the word $\sigma(a)$ ends with a letter from A_l. Clearly, if $A = A_l$ then σ has the nesting property.

The paper is organized as follows:

Section 2: Bratteli-Vershik models of Cantor aperiodic systems. In the section, we consider aperiodic homeomorphisms of a Cantor set and discuss the notions of an ordered Bratteli diagram and the Vershik map associated to aperiodic Cantor systems. We outline the proof of the fact that any aperiodic homeomorphism of a Cantor set is conjugate to the Vershik map of an ordered Bratteli diagram (see [Med]). This result is a foundation stone for our further research.

Section 3: Downarowicz-Maass' symbolic representation. In the paper [DM] the authors suggested a method of coding of dynamics on Bratteli diagrams by means of *j-symbols*. In the section, we generalize ideas and results from [DM] and give abstract definitions of j-symbols, j-sequences, etc. They can be used to study dynamics of different nature, for instance, Bratteli-Vershik systems and substitutional dynamical systems. The main advantage of this approach is that it allows one to use the machinery of symbolic dynamics for solving some problems of Cantor dynamics. The technique used in this section (see Propositions 3.10 and 3.12) is applied in the proofs of our main results.

Section 4: Finite rank aperiodic Cantor systems. We apply in this section the technique of j-sequences developed in Section 3 to the study of Cantor aperiodic systems of finite rank. Generalizing the main result of [DM], we show that the Vershik map of an ordered Bratteli diagram with uniformly bounded number of vertices at each level whose restrictions to minimal components are not conjugate to odometers is expansive, i.e. this map is homeomorphic to a subshift over a finite alphabet.

Section 5: Recognizability of aperiodic substitutional systems. We discuss in this section the properties of an arbitrary substitution σ over a finite alphabet A and the substitutional dynamical system (X_σ, T_σ) defined by σ (the rigorous definitions are given in Section 5). We prove that the number of minimal components of the system (X_σ, T_σ) cannot exceed $|A|$. We also show that for an arbitrary substitution σ any point $x \in X_\sigma$ can be written as

$$x = T_\sigma^i \sigma(y) \text{ for some } y \in X_\sigma \text{ and some } i = 0, \ldots, |\sigma(y[0])| - 1. \qquad (1)$$

If σ is a primitive aperiodic substitution, then representation (1) is unique for each x. This fact is a consequence of recognizability property of primitive substitutions established by Mossé [Mos1, Mos2]. We generalize this result by showing that if we omit the condition of primitivity for σ, then the uniqueness of the representation (1) still holds. The proof of the result involves the ideas of Downarowicz-Maass' symbolic representation [DM] and the technique developed in Section 3.

Section 6: Stationary Bratteli-Vershik models vs. aperiodic substitutional systems. In the section we show that every expansive Vershik map of a stationary ordered Bratteli diagram is homeomorphic to a substitutional dynamical systems. Conversely, we show that every aperiodic substitutional dynamical system constructed by a substitution satisfying the *nested* property is homeomorphic to the Vershik map of a stationary ordered Bratteli diagram. These results generalize those proved in [DHS] for minimal (primitive) substitutional systems.

Section 7: Derivative substitutions. The section is devoted to the study of derivative substitutions associated with aperiodic substitutions. This notion was first considered in [DHS] and [Dur] for primitive substitutions. We show how using derivative substitutions one can find a Bratteli-Vershik realization of a substitutional system. This approach differs from that of Section 6 and sometimes gives a 'simpler' Bratteli diagram.

Appendix: Description of the phase space X_σ. In the appendix we give an explicit description of elements from the space X_σ and establish some supplementary results on general substitutional systems.

2 Bratteli-Vershik Models of Cantor Aperiodic Systems

In this section we show how one can associate a Bratteli-Vershik dynamical system to every Cantor aperiodic (non-minimal) homeomorphism. This result was announced in [Med]. Since the notion of Bratteli diagrams has been discussed in many well known papers on Cantor dynamics (e.g. [HPS], [GPS1], [GPS3]), we give only the main steps of the construction.

By a *Cantor set* X we understand any zero-dimensional compact metric space without isolated points. Recall that a homeomorphism T is called *aperiodic* if every T-orbit is infinite; a homeomorphism $T : X \to X$ is called *minimal* if every orbit of T is dense in X.

Definition 2.1. A *Bratteli diagram* is an infinite graph $B = (V, E)$ such that the vertex set $V = \bigcup_{i\geq 0} V_i$ and the edge set $E = \bigcup_{i\geq 1} E_i$ are partitioned into disjoint sets V_i and E_i such that
(i) $V_0 = \{v_0\}$ is a single point;
(ii) V_i and E_i are finite sets;
(iii) there exist a range map r and a source map s from E to V such that $r(E_i) = V_i$, $s(E_i) = V_{i-1}$, and $s^{-1}(v) \neq \emptyset$, $r^{-1}(v') \neq \emptyset$ for all $v \in V$ and $v' \in V \setminus V_0$.

The pair (V_i, E_i) is called the i-th level of the diagram B. We write $e(v, v')$ to denote an edge e such that $s(e) = v$ and $r(e) = v'$.

A finite or infinite sequence of edges $(e_i : e_i \in E_i)$ such that $r(e_{i-1}) = s(e_i)$ is called a finite or infinite path, respectively. It follows from the definition that every vertex $v \in V$ is connected to v_0 by a finite path and the set $E(v_0, v)$ of all such paths is finite. For a Bratteli diagram B, we denote by X_B the set of infinite paths. We endow the set X_B with the topology generated by cylinder sets $U(e_1, \ldots, e_n) = \{x \in X_B : x_i = e_i,\ i = 1, \ldots, n\}$, where $(e_1, \ldots, e_n)$ is a finite path of B. The set X_B is a 0-dimensional compact metric space with respect to this topology.

Remark 2.2. In general, the set X_B may have isolated points. We do not assume that X_B is a perfect space; our results remain true for any 0-dimensional compact metric space.

Let $B = (V, E, \leq)$ be a Bratteli diagram (V, E) equipped with a partial order $\leq$ defined on each $E_i,\ i = 1, 2, ...$, such that edges e, e' are comparable if and only if $r(e) = r(e')$; in other words, a linear order $\leq$ is defined on each (finite) set $r^{-1}(v),\ v \in V \setminus V_0$. For a Bratteli diagram (V, E) equipped with such a partial order $\leq$ on E, one can also define a partial lexicographic order on the set $E_{k+1} \circ \cdots \circ E_l$ of all paths from V_k to V_l: $(e_{k+1}, ..., e_l) < (f_{k+1}, ..., f_l)$ if and only if for some i with $k + 1 \leq i \leq l$, $e_j = f_j$ for $i < j \leq l$ and $e_i < f_i$. Then any two paths from $E(v_0, v)$ are comparable with respect to the introduced lexicographic order. We call a path $e = (e_1, e_2, ..., e_i, ...)$ *maximal (minimal)* if every e_i is maximal (minimal) amongst all elements from $r^{-1}(r(e_i))$. Notice that there are unique minimal and maximal paths in $E(v_0, v)$ for each $v \in V_i,\ i \geq 0$.

Denote the sets of all maximal and minimal paths in X_B by X_{max} and X_{min}, respectively. It is not hard to see that X_{max} and X_{min} are non-empty closed sets.

Definition 2.3. A Bratteli diagram $B = (V, E)$ together with a partial order $\leq$ on E is called *an ordered Bratteli diagram* $B = (V, E, \leq)$.

Definition 2.4. Let $B = (V, E, \leq)$ be an ordered Bratteli diagram. We say that a homeomorphism $\varphi_B : X_B \to X_B$ is a *Vershik homeomorphism (map)* if it satisfies the following conditions:
(i) $\varphi_B(X_{max}) = X_{min}$;
(ii) if $x = (x_1, x_2, \ldots) \notin X_{max}$, then

$$\varphi_B(x_1, x_2, \ldots) = (x_1^0, \ldots, x_{k-1}^0, \overline{x_k}, x_{k+1}, x_{k+2}, \ldots),$$

where $k = \min\{n \geq 1 : x_n \text{ is not maximal}\}$, $\overline{x_k}$ is the successor of x_k in the set $r^{-1}(r(x_k))$, and $(x_1^0, \ldots, x_{k-1}^0)$ is the minimal path in $E(v_0, s(\overline{x_k}))$.

It is well-known that every simple Bratteli diagram B has an ordering such that the sets $X_{\max}$ and $X_{\min}$ are singletons. In contrast to this case, non-simple Bratteli diagrams may not have such an ordering, in general. We notice also that not every order of a Bratteli diagram defines a continuous Vershik map φ_B. Moreover, it was shown in [Med] that *there exists a Bratteli diagram B such that any ordering of B does not produce the (continuous) Vershik map.*

Let $B = (V, E, \leq)$ be an ordered Bratteli diagram such that the Vershik map φ_B exists. Then the pair (X_B, φ_B) is called a *Bratteli-Vershik system.*

To construct a Bratteli diagram associated to a homeomorphism of a Cantor set, one needs to work with sequences of Kakutani-Rokhlin partitions. For a minimal homeomorphism, such sequences can be easily produced via the first return functions for any clopen sets.

We consider now aperiodic Cantor systems and show how we can construct Bratteli diagrams in this case [BDM], [Med].

Definition 2.5. Let (X, T) be a Cantor dynamical system. (1) By a *Kakutani-Rokhlin (K-R) partition* $\mathcal{P}$ of X, we understand any partition of X into clopen sets of the form

$$\mathcal{P} = \{T^i A(v)\ :\ v = 1, \ldots, m;\quad i = 0, \ldots, h(v) - 1\}.$$

The *base* of the partition $\mathcal{P}$ is $B(\mathcal{P}) = \bigcup_{v=1}^{m} A(v)$. A family of disjoint sets $\xi(v) = \{A(v), TA(v), \ldots, T^{n-1}A(v)\}$ is called a *T-tower* of *height* $h(\xi) = n$. Then $\mathcal{P}$ is the disjoint union of clopen T-towers $\xi(v)$.

(2) A sequence of K-R partitions $\{\mathcal{P}_n\}$ is called *nested* if $\mathcal{P}_{n+1}$ refines $\mathcal{P}_n$ and $B(\mathcal{P}_{n+1}) \subseteq B(\mathcal{P}_n)$ for any n.

For a Cantor aperiodic system (X, T), a clopen set A is called a *complete T-section* if A meets every T-orbit at least once. A point $x \in A$ is called *recurrent* with respect to A if there exists $n \in \mathbb{N}$ such that $T^n x \in A$. By compactness of X, every clopen complete T-section A consists of recurrent points. Thus, using the first return function $n_A(x) = \min\{n \in \mathbb{N} : T^n x \in A\}$ for a complete T-section A, we can construct a K-R partition $\mathcal{P}$ of X with $B(\mathcal{P}) = A$.

Theorem 2.6 *[Med] Let (X, T) be a Cantor aperiodic system. There exists a sequence of K-R partitions $\{\mathcal{P}_n\}$ of X such that for all $n \geq 1$: (i) $\mathcal{P}_{n+1}$ refines $\mathcal{P}_n$; (ii) $h_{n+1} > h_n$, where $h_n = \min\{h(\xi) : \xi \in \mathcal{P}_n\}$; (iii) $B(\mathcal{P}_{n+1}) \subseteq B(\mathcal{P}_n)$; (iv) the elements of partitions $\{\mathcal{P}_n\}$ generate the clopen topology on X.*

Let (X, T) be a Cantor aperiodic system. We say that a closed set $Y \subset X$ is a *basic* set if every clopen neighborhood of Y is a complete T-section and Y meets every T-orbit at most once. If a sequence of K-R partitions $\{\mathcal{P}_n\}$ satisfies the conditions of Theorem 2.6, then $Y = \bigcap_n B(\mathcal{P}_n)$ is a basic set.

Theorem 2.7 *[Med] Let (X, T) be a Cantor aperiodic system with a basic set Y. There exists an ordered Bratteli diagram $B = (V, E, \leq)$ such that (X, T) is conjugate to a Bratteli-Vershik model (X_B, φ_B). The homeomorphism implementing the conjugacy between T and φ_B maps the basic set Y onto the set $X_{\min}$ of all minimal paths of X_B.*

Our primary goal is to construct Bratteli-Vershik models for aperiodic substitutional systems. We recall briefly the main steps of this construction for an aperiodic homeomorphisms of a Cantor set.

Let (X, T) be a Cantor aperiodic system. By Theorem 2.6 find a sequence of nested K-R partitions: $\mathcal{P}_0 = X$,

$$\mathcal{P}_n = \{T^i A(v, n) \ : \ v = 1, \ldots, m(n); \ \ i = 0, \ldots, h(v, n) - 1\}, \ n \geq 1, \tag{2}$$

that generates the clopen topology on X. Set

$$\xi(v, n) = \{A(v, n), \ldots, T^{h(v,n)-1}A(v, n)\} \text{ for } v = 1, \ldots, m(n).$$

Define an ordered Bratteli diagram $B = (V, E, \leq)$ as follows:

(i) Define the set of vertices by $V_0 = \{v_0\}$ and $V_n = \{1, \ldots, m(n)\}$ for $n \geq 1$.

(ii) Define the set of edges E_n between the consecutive levels V_{n-1} and V_n by the incidence matrix $M(n) = \{m_{vw}(n) : v \in V_n, \ w \in V_{n-1}\}$, where

$$m_{vw}(n) = |\{0 \leq i < h(v, n) : T^i A(v, n) \subset A(w, n-1)\}|.$$

In other words, we fix a vertex $v \in V_n$ and define $V(v, n-1)$ as the set of all vertices from V_{n-1} such that $\xi(v,n)$ intersects $\xi(w, n-1)$, and a vertex w appears in $V(v, n-1)$ as many times as $\xi(v,n)$ intersects $\xi(w, n-1)$. Then we connect v to each vertex $w \in V(v, n-1)$ taking into account the multiplicity of appearance of w in $V(v, n-1)$.

(iii) To define the ordering $\leq$ on E we take the clopen set $A(v,n)$. Then tracing the orbit of $A(v,n)$ within the T-tower $\xi(v,n)$, we see that $T^i A(v,n)$ consecutively meets the sets $A(w_1, n-1)$, $A(w_2, n-1)$,..., $A(w_{k_v}, n-1)$ (some of them can occur several times). This defines the set of edges $r^{-1}(v)$. Enumerate the edges from $r^{-1}(v)$ as follows: $e(w_1, v) < e(w_2, v) < \ldots < e(w_{k_v}, v)$.

Since the partitions $\{\mathcal{P}_n\}$ generate the topology of X, for each point $x \in X$ there is a unique sequence $i(x) = \{(v_n, i_n)\}_{v \in V_n;\, 0 \leq i_n < h(v_n, n)}$ such that

$$\{x\} = \bigcap_{n=1}^{\infty} T^{i_n} A(v_n, n).$$

Define the map $\theta : X \to X_B$ by

$$\theta(x) = \bigcap_{n \geq 1} U(y_1, \ldots, y_n)$$

where $(y_1, \ldots, y_n)$ is the i_n-th finite path in $E(v_0, v_n)$ with respect to the lexicographical ordering on $E(v_0, v_n)$. It can be easily checked that θ is a homeomorphism.

Define $\varphi_B = \theta \circ T \circ \theta^{-1}$. Clearly, the homeomorphism φ_B satisfies Definition 2.4 with the ordering $\leq$. Thus, we obtain that (X, T) is conjugate to the Bratteli-Vershik system (X_B, φ_B).

Open question. It is well-known that for every simple properly ordered Bratteli diagram $B = (V, E)$ there exists the continuous Vershik map, see [HPS]. On the other hand there are Bratteli diagrams such that any ordering on them will not lead to a *continuous* Vershik map [Med]. The question is: How can one describe the class of Bratteli diagrams which admit continuous Vershik maps on the set of infinite paths?

3 Downarowicz-Maass' Symbolic Representation

In the section, we develop the ideas of the paper [DM] and apply them below to solving the following problems: recognizability of aperiodic substitutions (Section 5) and expansiveness of Vershik maps (Section 4). The results stated in this section are proved in [DM]. Since they are of crucial importance for us, we reproduce their proofs here with minor alterations.

Fix a sequence of finite alphabets $\{A_i\}_{i \geq 0}$.

Definition 3.1. (1) Take $a \in A_j$, $j \geq 1$. By a *j-symbol* $[a]_j$ we understand a finite matrix with $j+1$ rows numbered from $\{0, 1, \ldots, j\}$ which has the following structure:

(i) The row j (the bottom row of $[a]_j$) consists of one box carrying the symbol a (in other words, labeled by a) whose length is extending over the full width of the matrix.

(ii) The row $j-1$ consists of m_{j-1} boxes labeled by letters $b_0, \dots, b_{m_{j-1}-1}$ from the alphabet A_{j-1}. The total length of these boxes is equal to the width of the matrix $[a]_j$.

(iii) The row $j-2$ consists of m_{j-1} group of boxes. The i-th group, $i = 0, \dots, m_{j-1}-1$, is located exactly over the box of $(j-1)$-th row labeled by b_i; the total length of boxes from the group i is equal to the length of the box b_i. Each box from the row $j-2$ is labeled by a letter from the alphabet A_{j-2}.

(iv) All rows above have the same structure. The row 0 (the first row of the matrix) consists of m_0 boxes of length 1 which are labeled by letters from A_0.

(2) A *0-symbol* is always a 1×1 matrix viewed as a box of length 1 which is labeled by a letter from A_0.

We observe that the set of all j-symbols, $j \geq 1$, is infinite. Denote by $\mathcal{A}_j$ any finite set of j-symbols over the alphabets $A_0, \cdots, A_j$, $j \geq 0$. When we use the notation $\mathcal{A}_j$, it will be clear from the context which set of j-symbols we mean.

Definition 3.2. Let maps $\sigma_i : A_i \to A_{i-1}^+$, $i \geq 1$, be given. For every i define by induction the family of i-symbols $\{\mathcal{A}_i\}_{i \geq 0}$:

(1) $\mathcal{A}_0 = \{[a]_0 : a \in A_0\}$.

(2) Suppose that the family $\mathcal{A}_{i-1}$ is defined. For $a \in A_i$, let $\sigma_i(a) = a_0 \cdots a_m$ where $a_0, \dots, a_m \in A_{i-1}$. Define the i-symbol $[a]_i$ as the matrix whose first i rows are concatenations of $(i-1)$-symbols $[a_0]_{i-1}, \dots, [a_m]_{i-1}$ from $\mathcal{A}_{i-1}$. Then the family $\mathcal{A}_i$ is formed by i-symbols $\{[a]_i : a \in A_i\}$.

We call this families of i-symbols $\{\mathcal{A}_i\}_{i \geq 0}$ *agreeable.*

Example 3.3. Let $A_2 = \{a, b, ...\}$, $A_1 = \{x, y...\}$, $A_0 = \{\alpha, \beta, ...\}$. Then the following matrix gives an example of a 2-symbol $[a]_2$:

<table>
<tr><td>α</td><td>β</td><td>γ</td><td>δ</td></tr>
<tr><td colspan="2">x</td><td colspan="2">y</td></tr>
<tr><td colspan="4">a</td></tr>
</table>

In this example, we have that $\sigma_2(a) = xy$, $\sigma_1(x) = \alpha\beta$, and $\sigma_1(y) = \gamma\delta$ where $\sigma_i : A_i \to A_{i-1}^+$, $i = 1, 2$.

Remark 3.4. Suppose that the families of j-symbols $\{\mathcal{A}_j\}_{j \geq 0}$ are agreeable. Then any row $i \leq j$ of a j-symbol $[a]_j \in \mathcal{A}_j$ completely determines all rows above. Furthermore, it follows from Definition 3.2 that each letter $a \in A_j$ is uniquely assigned to the j-symbol $[a]_j$ from $\mathcal{A}_j$.

We give now two principal constructions of agreeable families.

Example 3.5. Let $B = (V, E, \leq)$ be an ordered Bratteli diagram with the path space X_B. Set $A_i = V_i$, $i \geq 0$. Define the map $\sigma_i : A_i \to A_{i-1}^+$ $(i \geq 1)$

as follows: take a vertex $v \in V_i$, write down all edges $e_1 < e_2 < \ldots < e_m$ from $r^{-1}(v)$ with respect to the ordering $\leq$, and define

$$\sigma_i(v) = v_1 \ldots v_m$$

where $v_k = s(e_k)$, $k = 1, \ldots, m$. Then the alphabets $\{A_i\}_{i \geq 0}$ and maps $\sigma_i : A_i \to A_{i-1}^+$ define the sets $\{\mathcal{A}_j\}_{j \geq 0}$ of agreeable j-symbols, see Definition 3.2.

To illustrate this example consider the following Bratteli diagram with the edges ordered from the left to right.



Then the 2-symbol $[c]_2$ can be viewed as the matrix

<table>
<tr><td>v_0</td><td>v_0</td><td>v_0</td><td>v_0</td></tr>
<tr><td>a</td><td>b</td><td>c</td><td>c</td></tr>
<tr><td colspan="4">c</td></tr>
</table>

Example 3.6. Let $\tau : A \to A^+$ be a substitution defined on a finite alphabet A. Set $A_0 = A$ and $A_j = \{\tau^j(a) : a \in A\}, j > 0$. Define $\sigma_j : A_j \to A_{j-1}^+$ by setting $\sigma_j(\tau^j(a)) = \tau^{j-1}(a_1) \cdots \tau^{j-1}(a_m)$ where $a_1 \cdots a_m = \tau(a)$. It follows from Definition 3.2 that the data $\{(A_j, \sigma_j)\}_{j \geq 0}$ define the agreeable families of j-symbols $\{\mathcal{A}_j\}_{j \geq 0}$.

Definition 3.7. Let $\{\mathcal{A}_0, \ldots, \mathcal{A}_j\}$ be agreeable families of symbols. Denote by $\mathcal{Z}_j$ the set of all matrices with $j+1$ two-sided infinite rows which are obtained by concatenation of j-symbols. Every element of $\mathcal{Z}_j$ is a sequence $\{[a_n]_j\}_{n \in \mathbb{Z}}$ with $[a_n]_j \in \mathcal{A}_j$. Elements of $\mathcal{Z}_j$ are called *j-sequences.*

For $x \in \mathcal{Z}_j$ and $n \in \mathbb{Z}$, denote by $x(n)$ the n-th column of x. The i-th row of the matrix x $(i < j)$ is a concatenation of boxes of variable lengths which are labeled by i-symbols from $\mathcal{A}_i$. So it is natural to represent $x(n)$ as a column whose i-th entry carries the following information: the letter $a \in A_i$ such that the i-symbol $[a]_i$ intersects the column $x(n)$ in x and an integer $k > 0$ that denotes the coordinate of the column $x(n)$ within the i-symbol $[a]_i$.

This observation allows us to write down the matrix (j-sequence) x as the sequence $x = \{x(n)\}_{n \in \mathbb{Z}}$. Define $T : \mathcal{Z}_j \to \mathcal{Z}_j$ by $(Tx)(n) = x(n+1)$ for all $n \in \mathbb{Z}$. Then $(\mathcal{Z}_j, T)$ is a subshift over a finite alphabet.

The following picture gives an example of a 2-sequence.

<table>
<tr><td>...</td><td>x_{-6}</td><td>x_{-5}</td><td>x_{-4}</td><td>x_{-3}</td><td>x_{-2}</td><td>x_{-1}</td><td>x_0</td><td>x_1</td><td>x_2</td><td>x_3</td><td>x_4</td><td>x_5</td><td>x_6</td><td>...</td></tr>
<tr><td>...</td><td colspan="3">y_{-2}</td><td colspan="2">y_{-1}</td><td colspan="4">y_0</td><td colspan="4">y_1</td><td>...</td></tr>
<tr><td>...</td><td colspan="5">z_{-1}</td><td colspan="8">z_0</td><td>...</td></tr>
</table>

Definition 3.8. Let $\{\mathcal{A}_0, \mathcal{A}_1, \ldots\}$ be agreeable families of symbols and $x, y \in \mathcal{Z}_j$ for some $j > 0$.

(1) x and y are called *i-compatible*, $i \leq j$, if the i-th rows of x and y coincide. Observe that since the families $\{A_k\}_{k \geq 0}$ are agreeable, the i'-th rows of x and y also coincide for all $i' < i$.

(2) If $x \neq y$ and they are compatible, then the maximal $i \leq j$ such that the i-th rows of x and y coincide is called the *depth* of x and y.

(3) If the i-th rows of x and y are different, then x and y are called *i-separated.*

(4) x and y are said to have a *common j-cut* if there exist $n \in \mathbb{N}$ and j-symbols $[a]_j$ and $[b]_j$ such that $[a]_j$ and $[b]_j$ appear in x and y, respectively, at the position n.

Remark 3.9. (1) We note that if a pair (x, y) has a common j-cut at a coordinate n, then it has common i-cuts for all $i \leq j$ at the same coordinate n.

(2) For a j-symbol $[a]_j$, let $|[a]_j|$ stand for the width of the matrix $[a]_j$. We observe that if a j-symbol $[a]_j$ appears in $x \in \mathcal{Z}_j$ at the position n, then there is a j-symbol $[a']_j$ that appears in x at the position $n + |[a]_j|$.

(3) Denote by $\pi_i : \mathcal{Z}_j \to \mathcal{Z}_i$, $i < j$, the map that restricts each $x \in \mathcal{Z}_j$ to its first $(i+1)$ rows. Then π_i is a factor map from $(\mathcal{Z}_j, T)$ onto $(\mathcal{Z}_i, T)$.

It is not hard to see that the technique of j-symbols is an interpretation of the well-known technique of Kakutani-Rokhlin partitions. However, the usage of j-symbols sometimes is more convenient as it allows us to manipulate with blocks, symbols, and other symbolic objects.

Proposition 3.10 *[DM] Let $\{\mathcal{A}_j\}_{j \geq 0}$ be agreeable families of j-symbols. Suppose also that $|\mathcal{A}_j| \leq K < \infty$ for all $j \geq 0$. Then for any $n \in \mathbb{N}$ there exist $i, j \in \mathbb{N}$ with $n \leq i < j$ such that any pair of j-sequences with depth i has no common j-cuts.*

Proof. Assume the converse. Then take $n_0 \in \mathbb{N}$ such that for any $i = n_0, \ldots, j-1$ there exists a pair (x_i, y_i) of j-sequences with depth i that has a common j-cut. We set $j = n_0 + K$.

(I) Consider the pair (x_{j-1}, y_{j-1}). By assumption, this pair is $(j-1)$-compatible, j-separated and has a common j-cut.

Fix any common j-cut for x_{j-1} and y_{j-1}, i.e. we fix a position n at which possibly different j-symbols appear in x_{j-1} and y_{j-1}. Consider the j-symbols occurring at the position n in x_{j-1} and y_{j-1}. If these j-symbols are the same, then the following j-cut is also common and we consider the following j-symbols (to the right). Let u and v be the first different j-symbols to the right of n. If there are no such symbols, we analyze x_{j-1} and y_{j-1} to the left of n.

If the j-symbols u and v have the same length, then the following j-cut is also common. In this case, we do not change anything and continue checking the following pair of j-symbols.

If, say, u is longer than v, we modify the set of j-symbols. We replace the j-symbol u from $\mathcal{A}_j$ by the concatenation of two j-symbols, $v \in \mathcal{A}_j$ and a new symbol u' which is defined as follows. The matrix $[u']_j$ is formed by $|u| - |v|$ the right most columns of the matrix $[u]_j$ with the $(j+1)$-th row labeled by u'. In other words, we substitute the last row in the symbol u by two boxes: v and a new symbol u' which has the complementary length. Observe that the rows above the last one are not changed (recall that x_{j-1} and y_{j-1} are $(j-1)$-compatible).

Next we replace occurrence of the j-symbol u in every element of $\mathcal{Z}_j$ by the concatenation vu'. This procedure lead to a symbolic system which is topologically conjugate to $(\mathcal{Z}_j, T)$. Note also that we produce more j-cuts and never remove them. If there is still a non-common cut, we repeat the described construction and substitute some j-symbol with an existing one and a new one.

As soon as we get that all j-cuts to the right of n are common, we repeat the argument to the left part of x_{j-1} and y_{j-1}.

(II) Thus, we get a symbolic system topologically conjugate to $\mathcal{Z}_j$ with the same number of j-symbols such that the modified sequences x_{j-1} and y_{j-1} have all j-cuts common. On the other hand, the sequences x_{j-1} and y_{j-1} remain $(j-1)$-compatible as we did not change the upper j-rows. Furthermore, x_{j-1} and y_{j-1} remain j-separated. The latter means that there are two different j-symbols which are $(j-1)$-compatible.

Now we produce a topological factor $\mathcal{Z}'_j$ of $\mathcal{Z}_j$ by identifying j-symbols which are $(j-1)$-compatible. Thereby, we strictly reduce the number of j-symbols and j-sequences from $\mathcal{Z}'_j$ have at most $K-1$ different j-symbols.

(III) It follows from (II) that the sequences x_{j-1} and y_{j-1} are not j-separated any more. The every pair (x_i, y_i), $i \in [n_0, j-2]$ (in fact, their images in the factor $\mathcal{Z}'_j$) remains $(i+1)$-separated since these rows have not been changed. Moreover, the pair (x_i, y_i) still has a common j-cut because in the construction we only added more cuts.

We can now apply the same arguments as in (I) and (II) to the pair (x_{j-2}, y_{j-2}). As a result, we get a new factor with at most $K-2$ j-symbols.

Repeating the same argument no more than $K-1$ times, we obtain a new factor in which the pair (x_{n_0}, y_{n_0}) remains (n_0+1)-separated with a common j-cut while the family of j-symbols consists of one element only. This is a contradiction. $\square$

The next proposition is called the "Infection lemma" in [DM]. Before proving the result, we recall the notion of an eventually periodic sequence.

Definition 3.11. A sequence $x = \{x(n)\}_{n\in\mathbb{Z}}$ is called *eventually periodic* if there are n_0 and m such that $x(n+m) = x(n)$ for all $n \geq n_0$.

Recall also that the map $\pi_i : \mathcal{Z}_j \to \mathcal{Z}_i$ denotes the projection to the first $i+1$ lines.

Proposition 3.12 *[DM] Let $\{\mathcal{A}_0, \dots, \mathcal{A}_j\}$ be agreeable families of j-symbols with $|\mathcal{A}_i| \leq K$, $i = 0, \dots, j$, where $j \geq K^2$. Suppose that the set $\{z_0, \dots, z_{K^2}\}$ consists of i-compatible ($i < j$) and pairwise j-separated j-sequences with no common j-cuts. Then $\pi_i(z_0)$ is eventually periodic.*

Proof. (a) We can set $\hat{z} := \pi_i(z_k)$ because all the points z_k are i-compatible. Analogously, denote by $\hat{v}$ the restriction of a j-symbol v to its top $i+1$ rows, i.e. the projection of v to $\mathcal{A}_i$. Draw a diagram $\mathcal{D}$ consisting of the j-th rows of the elements z_k one above another with aligned zero coordinate.



Diagram $\mathcal{D}$: j-th rows without common j-cuts.

(b) Since we have K^2+1 sequences of j-symbols $\{z_0, \dots, z_{K^2}\}$, zero coordinates of $\{z_0, \dots, z_{K^2}\}$ are covered by at least $K+1$ copies of a j-symbol v. Notice that since the elements $\{z_0, \dots, z_{K^2}\}$ have no common j-cuts, any of two copies of v are shifted by some positive integer $0 < l < |v|$. Using the fact that $\pi_i(z_k) = \pi_i(z_0)$, we get that the projection of the j-symbol v to $\mathcal{A}_i$ satisfies the "l-periodicity law": $\hat{v}(n) = \hat{v}(n+l)$ for every $n \in [0, |v|-1-l]$.

(c) Let l_v be the minimal shift for v appearing in the diagram $\mathcal{D}$. This means that $\hat{v}(n) = \hat{v}(n+l_v)$ for every $n \in [0, |v|-1-l_v]$. As zero coordinate of $\hat{z}$ is covered by at least two copies of v, we have that $\hat{z}(0) = \hat{z}(l_v)$. Denote by I the largest interval of $\mathbb{Z}$ such that $0 \in I$ and if $n \in I$, then $\hat{z}(n) = \hat{z}(n+l_v)$.

(d) If I is not bounded to the right, then the sequence $\hat{z} = \pi_i(z_0)$ is eventually periodic and we are done.

If I has the right end, set $m = (\max_{n \in I} n) + 1$. Hence $\hat{z}(m) \neq \hat{z}(m+l_v)$. Restrict the diagram $\mathcal{D}$ to those $K+1$ elements z_k in which the coordinate 0 is covered by the j-symbol v. Since this diagram consists of $K+1$ lines, the coordinate m is covered by at least two copies of a j-symbol w.

If $w = v$, then by the choice of l_v, we get that $\hat{z}(m) = \hat{z}(m+l_v)$, which is impossible.

Suppose that $w \neq v$. Let r be the relative coordinate of m within the extreme left copy of w covering m (see the Figure[1] below). Since there is no common j-cuts, $r > 0$. Hence, absolute coordinates of the extreme left copy of $\hat{w}[0, r-1]$ within $\hat{z}$ intersects I. As $w \neq v$, we have that the absolute coordinates of $\hat{w}[0, r-1]$ lie in I, which implies that $\hat{w}(n) = \hat{w}(n+l_v)$ for all $n \in [0, r-1]$.

---

[1] The idea of the figure is taken from [DM].


I
$\dots$ $\hat{z}_{-2}$ $\hat{z}_{-1}$ $\hat{z}_0$ $\hat{z}_1$ $\dots$ $\hat{z}_{m-r}$ $\dots$ $\hat{z}_{m-n}$ $\dots$ $\hat{z}_{m-1}$ $\hat{z}_m$ $\dots$ $\dots$ $\dots$
z_{k_1} v w_0 $\dots$ w_{r-n} $\dots$ w_{r-1} w_r $\dots$
z_{k_2} v w_0 $\dots$ w_{n-1} w_n $\dots$ $\dots$ $\dots$

Now, consider the other copy of w. Since it is the extreme right copy of w, the position m is aligned with some relative position $n \in [0, r-1]$, where the l_v-periodicity law holds. Thus, $\hat{z}(m) = \hat{z}(m + l_v)$, which is a contradiction. $\square$

4 Finite Rank Systems

In the section we generalize the result of [DM] to homeomorphisms of a Cantor set with finite rank. To prove our main result of this section we develop the method used in [DM] to the case of aperiodic homeomorphisms.

Let $B = (V, E, \leq)$ be an ordered Bratteli diagram such that the ordering $\leq$ admits the continuous Vershik map $\varphi_B : X_B \to X_B$ where X_B is the space of infinite paths. Denote by $X_{\max}$ and $X_{\min}$ the sets of all maximal and minimal paths of X_B, respectively. Recall that by definition of the Vershik map we have that $\varphi_B(X_{\max}) = X_{\min}$.

Let $\{\mathcal{A}_j\}_{j \geq 0}$ be the family of agreeable j-symbols associated to the diagram B (see the details in Example 3.5).

Remark 4.1. (1) Any infinite path $x = (x_n) \in X_B \setminus Orb_{\varphi_B}(X_{min} \cup X_{max})$ can be represented as an infinite matrix denoted by $[x]$ with rows indexed by i from 0 to ∞ and columns indexed by $j \in (-\infty, \infty)$. The matrix $[x]$ is formed by an increasing sequence of i-symbols ($i = 0, 1, \ldots$) which cross the coordinate 0. The i-symbol corresponds to the vertex $v \in V_i$ which is crossed by the path x at the level i, and the position of 0 coordinate in this i-symbol is defined by the order of the finite path $(x_1, \ldots, x_i)$ amongst all paths connecting v_0 and v. We see that this construction gives not only a single point $x \in X_B$ but the entire orbit $Orb_{\varphi_B}(x) = \{\varphi_B^i(x)\}_i$.

(2) We notice that for every $x \in X_{\max}$ the Vershik map uniquely defines $y \in X_{\min}$ such that $y = \varphi_B(x)$. If we applied the construction used in (1) to the paths x and y, we would get two one-sided matrices $[y]^+$ and $[x]^-$ infinite to the right and left, respectively. As $Orb_{\varphi_B}(x) = Orb_{\varphi_B}(y)$ it is natural to assign the concatenated matrix $[x]^-[y]^+$ to x (or y).

(3) Let X be the set of infinite matrices described in (1) and (2). Then the Bratteli-Vershik model (X_D, φ_B) is conjugate to (X, T) where T is the left shift in X. This fact allows us to identify the sets X and X_B.

Recall that $\mathcal{Z}_i$ denotes the set of all i-sequences, see Definition 3.7. Let $\pi_i : X_B \to \mathcal{Z}_i$ be the map which restricts each matrix $[x]$, $x \in X_B$, to the first

$i+1$ rows. Set $X_i = \pi_i(X_B)$. Clearly, X_i is a closed shift-invariant subset of $\mathcal{Z}_i$. Observe also that (X_i, T) is a factor of (X_B, φ_B).

Definition 4.2. Let d be a metric on X which generates the topology. It is said that a homeomorphism $S : X \to X$ is *expansive* if there exists $\delta > 0$ such that for any distinct $x, y \in X$ there is $m \in \mathbb{Z}$ with $d(S^m x, S^m y) > \delta$. The number δ is called an *expansive constant*.

Note that the notion of expansiveness does not depend on the choice of the metric d, see [Wal, Section 5.6]

Remark 4.3. If (X_B, φ_B) is an expansive system, then, due to the famous theorem of Hedlund, (X_B, φ_B) is homeomorphic to (X_i, T) for all sufficiently large i, for the details see the proof of Theorem 5.24 in [Wal].

The following definitions agree with Definition 3.8.

Definition 4.4. We say that two distinct points x and y from X_B are *i-compatible* if $\pi_i(x) = \pi_i(y)$. If $\pi_i(x) \neq \pi_i(y)$, then x and y are called *i-separated*. Clearly, any distinct points x and y are i-separated for some i. The largest integer i such that $\pi_i(x) = \pi_i(y)$ is called the *depth* of x and y. A pair (x, y) has a common *j-cut* if there exist $n \in \mathbb{Z}$ and j-symbols v and w such that v appears at the position n in $\pi_j(x)$ and w appears at n in $\pi_j(y)$. Notice that if x and y have a common j-cut, then x and y have a common j'-cut for all $j' \leq j$.

Definition 4.5. A Cantor dynamical system (X, S) has the *topological rank* $K > 0$ if it admits a Bratteli-Vershik model (X_B, φ_B) such that the number of vertices of the diagram B at each level is not greater than K and K is the least possible number of vertices for any Bratteli-Vershik realization.

Clearly, if a system (X, S) has the rank K, then, by an appropriate telescoping, we can assume that the diagram B has exactly K vertices at each level.

The next statement shows that the number of minimal components of a finite rank system is bounded (see also Proposition 5.6 for a similar result for substitutional systems).

Proposition 4.6 *Let (X, S) be a Cantor aperiodic dynamical system of a finite rank K. Then (X, S) has at most K minimal components.*

Proof. By definition, (X, S) can be realized as a Bratteli-Vershik model with at most K vertices at each level. Assume that there exist $(K+1)$-minimal components $Z_0, \ldots, Z_K$ for a homeomorphism S. Then for each level i there exist a vertex $v_i \in V_i$ and two paths x_i and y_i from X_B such that they belong to different minimal components from $Z_0, \ldots, Z_K$ and pass through v_i. It follows that there exist distinct n and m such that $0 \leq n, m \leq K$ and $x_i \in Z_n,\ y_i \in Z_m$ for infinitely many indexes $i \in I$.

Define the metric d on the space X_B as follows:

$$d(\{x_n\}, \{y_n\}) = \frac{1}{\min\{k : x_k \neq y_k\}}. \tag{3}$$

Let $i_0 \in I$ be chosen such that $\mathrm{dist}(Z_n, Z_m) > 1/i_0$. Notice that we can find a path $z \in Orb_S(x_{i_0})$ such that the first i_0 edges of z coincide with those of y_{i_0}. It follows that $\mathrm{dist}(z, Z_m) \leq 1/i_0$, which is a contradiction. $\square$

We recall the definition of the enveloping semigroup, see, for example, the book [Gla] for a wider coverage of the subject.

Definition 4.7. Let $S : X \to X$ be a homeomorphism. By definition, the *enveloping semigroup* $E = E(X, S)$ of the dynamical system (X, S) is the closure of the set $\{S^n : n \in \mathbb{Z}\}$ in X^X with respect to the topology of pointwise convergence.

The main result of the section is the following statement.

Theorem 4.8 *Let (X, S) be an aperiodic Cantor dynamical system of finite rank K. If the restriction of (X, S) to every minimal component is not homeomorphic to an odometer, then (X, S) is expansive.*

Proof. Without loss of generality, we can assume that X is the path-space of an ordered Bratteli diagram $B = (V, E, \leq)$ and S is the Vershik map defined by the ordering $\leq$.

Assume that the system (X, S) is not expansive. Then for each $i \geq 1$ there exists a pair (x_i, y_i) such that $d(S^m x_i, S^m y_i) \leq 1/i$ for all $m \in \mathbb{Z}$, where the metric is defined in Equation 3. This is equivalent to the fact that the pair (x_i, y_i) is i-compatible. Therefore, for infinitely many i there is a pair of depth i. By telescoping of B, can assume that for each $i \geq 1$ there is a pair of depth i.

Now we have two opposite statements:

(1) There exists i_0 such that for all $i \geq i_0$ and every $j \geq i$ there is a pair of depth i with a common j-cut.

(2) For all i_0 there exist $i_0 \leq i < j$ such that any pair of depth i has no common j-cuts.

It follows from Proposition 3.10 that statement (1) is never true. Thus, for all i_0 there exist $j > i \geq i_0$ such that any pair of depth i has no common j-cuts. Therefore, any pair of depth i has no common j'-cuts for $j' > j$.

After telescoping of the diagram, we can assume that any pair of depth i has no common $(i+1)$-cuts. Recall that by our assumption of non-expansiveness of S, there exists at least one pair of depth i.

Let $E = E(X, S)$ be the enveloping semigroup for (X, S). Therefore, $Ex = \{\gamma(x) : \gamma \in E\} = \overline{Orb_S(x)}$. It follows that Ex contains at least one minimal S-component. By Proposition 4.6 there exist at most K minimal S-components, $Z_0, \ldots, Z_{K-1}$. Fix any path $z_i \in Z_i$, $i = 0, \ldots, K-1$.

Fix any level i_0. For every $i \in [i_0, i_0 + K^2 + K]$ take a pair (x_i, y_i) of depth i without common $(i+1)$-cuts. For each pair (x_i, y_i) find an element $\gamma_i \in E$ such that $\gamma_i(x_i) = z_j$ for some z_j, $j = j(i)$.

(a) Observe that the pair $(\gamma_i(x_i), \gamma_i(y_i))$ has the same depth as (x_i, y_i) and has no common $(i+1)$-cuts. Indeed, γ_i is the pointwise limit of a sequence S^{n_k}. As each projection $\pi_m : X \to X_m$ is continuous, we get

$$\begin{aligned} &\pi_i(\gamma_i(x_i)) = \pi_i(\lim S^{n_k} x_i) = \lim \pi_i(S^{n_k} x_i) = \lim T^{n_k} \pi_i(x_i) = \\ &= \lim T^{n_k} \pi_i(y_i) = \lim \pi_i(S^{n_k} y_i) = \pi_i(\gamma_i(y_i)). \end{aligned}$$

This, in particular, implies that $(\gamma_i(x_i), \gamma_i(y_i))$ is i-compatible.

It remains to show only that $(\gamma_i(x_i), \gamma_i(y_i))$ has no common $(i+1)$-cuts. Indeed, suppose $\gamma_i(x_i)$ and $\gamma_i(y_i)$ have a common $(i+1)$-cut. Then $T^{n_k}\pi_{i+1}(x_i)$ and $T^{n_k}\pi_{i+1}(y_i)$ have a common $(i+1)$-cut for all k sufficiently large. This implies that the pair (x_i, y_i) has a common $(i+1)$-cut, a contradiction.

(b) Consider the pairs $(\gamma_i(x_i), \gamma_i(y_i))$, $i \in [i_0, i_0 + K^2 + K]$. As each $\gamma_i(x_i)$ is equal to some z_j, $j = j(i)$, and we have at most K of them, we can choose K^2 pairs $(\gamma_i(x_i), \gamma_i(y_i))$, $i \in I$, such that $\gamma_i(x_i)$'s are the same for all $i \in I$, $|I| = K^2$, say $\gamma_i(x_i) = z_{k_0} \in Z_{k_0}$. Observe that the pair $(\gamma_i(y_i), \gamma_j(y_j))$, $i, j \in I$, $i < j$ has the depth i and has no common j-cuts.

(c) Setting $z_i' = \gamma_i(y_i)$ for $i \in I$, we obtain a family $C = \{z_i' : i \in I\} \cup \{z_{k_0}\}$ of i_0-compatible and pairwise $(i_0 + K^2 + K)$-separated elements with no common $(i_0 + K^2 + K)$-cuts. Considering the projection $\pi_{i_0+K^2+K}$ of C and applying Proposition 3.12, we get that $\pi_{i_0}(z_{k_0})$ is eventually periodic, and, by minimality, $\pi_{i_0}(Z_0)$ is periodic.

Thus, we get that for each i, there is a minimal S-component Z_i such that $\pi_i(Z_i)$ is T-periodic. As we have at most K minimal S-components, there is a minimal component, say Z_{k_0}, such that $\pi_i(Z_{k_0})$ is periodic for infinitely many i. This implies that the restriction of S to Z_{k_0} is homeomorphic to an odometer. The theorem is proved. □

5 Recognizability of aperiodic substitutions

In the section we study dynamical properties of an arbitrary substitutional dynamical system (X_σ, T_σ). First of all, we estimate the number of minimal components of the system (X_σ, T_σ). Then we build a sequence of K-R partitions of X_σ using geometrical properties of the substitutional dynamical system.

Let A denote a finite alphabet and A^+ the set of all non-empty words over A. Set also $A^* = A^+ \cup \{\emptyset\}$. For a word $w = w_0 \dots w_{n-1}$ with $w_i \in A$ let $|w| = n$ stand for its length. For any two words $v, w \in A^+$, the symbol '$v \prec w$' means that v is a factor of w.

Definition 5.1. By a *substitution* we mean any map $\sigma : A \to A^+$.

Any map $\sigma : A \to A^+$ is extended to the map $\sigma : A^+ \to A^+$ by concatenation. We define the *language* $L(\sigma)$ of a substitution σ as the set of all words which appear as factors of $\sigma^n(a)$, $a \in A$, $n \geq 1$. By definition, we also set that $\sigma^0(a) = a$ for all $a \in A$.

For any substitution $\sigma : A \to A^+$, define

$$|\sigma| = \min_{a\in A} |\sigma(a)|, \qquad ||\sigma|| = \max_{a\in A} |\sigma(a)|. \tag{4}$$

Observe that the functions $n \mapsto |\sigma^n|$ and $n \mapsto ||\sigma^n||$ are not decreasing.

Definition 5.2. By a *substitutional dynamical system associated to a substitution* σ, we mean a pair (X_σ, T_σ), where

$$X_\sigma = \{x \in A^{\mathbb{Z}} : x[-n, n] \in L(\sigma) \text{ for any } n\}$$

and T_σ is the shift on $A^{\mathbb{Z}}$. We will denote the k-th coordinate of $x \in X_\sigma$ by $x[k]$ or by $x(k)$. For any words $v, w \in L(\sigma)$, set $[v.w] := \{x \in X_\sigma : x[-|v|, |w| - 1] = vw\}$.

Definition 5.3. A substitution $\sigma : A \to A^+$ is called *aperiodic* if the system (X_σ, T_σ) has no periodic points.

Remark 5.4. (1) In general, the set X_σ can be empty. To avoid trivialities, *we will always assume that the substitution σ is such that X_σ is an uncountable subset of $A^{\mathbb{Z}}$*. In particular, we always have that $||\sigma^n||$ tends to the infinity.

(2) If $L(\sigma^k) = L(\sigma)$ for some $k \geq 1$, then the substitutional dynamical systems associated to σ and σ^k coincide, i.e., $X_\sigma = X_{\sigma^k}$.

(3) The set X_σ is a 0-dimensional compact metrizable space whose topology is generated by the metric

$$d(\{x[n]\}, \{y[n]\}) = \sum_{n=-\infty}^{\infty} \frac{d'(x[n], y[n])}{2^{|n|}} \tag{5}$$

where d' is the discrete metric on A, i.e. $d'(a, b) = 1$ iff $a \neq b$. Notice that the set X_σ can have isolated points. We do not require X_σ to be perfect.

Let $\sigma : A \to A^+$ be a substitution. We denote by A_l the set of all letters $a \in A$ such that $|\sigma^n(a)| \to \infty$ as $n \to \infty$. Set also $A_s = A \setminus A_l$. Here the subindexes 's' and 'l' stand for 'short' and 'long', respectively. Observe that $\sigma(a) \in A_s^+$ for all $a \in A_s$.

To estimate the number of minimal components of an arbitrary substitutional system, we need the following proposition which asserts that the length of words formed by short letters is uniformly bounded. This result will be also applied to Bratteli diagram construction (Theorem 6.6).

Proposition 5.5 *Let $\sigma : A \to A^+$ be an aperiodic substitution. Then there exists $M > 0$ such that every word $W \in L(X_\sigma)$ with $|W| \geq M$ contains at least one letter from A_l.*

Proof. Assume the converse, i.e., for any $m > 0$ there is a word $W_m \in L(X_\sigma) \cap A_s^+$ with $|W_m| = m$.

(0) Since $|\sigma^n(a)| \le K < \infty$ for all $a \in A_s$ and some K, there are positive integers n_a and p_a such that $\sigma^{n_a}(a) = \sigma^{n_a+kp_a}(a)$ for all $k \ge 0$, $a \in A_s$. Set $n_0 = \max\{n_a : a \in A_s\}$. Then it follows that

$$\sigma^n(a) = \sigma^{n+kp}(a) \text{ for all } k \ge 0 \text{ and } a \in A_s$$

where $n \ge n_0$ and $p = \prod_{a \in A_s} p_a$. Setting $\tau = \sigma^{pn_0}$, we obtain that $\tau^2(a) = \sigma^{pn_0+pn_0}(a) = \tau(a)$ for all $a \in A_s$, i.e. $\tau^k(a) = \tau(a)$ for all $a \in A_s$ and $k \ge 1$.

(1) Since $|A| < \infty$, we find a letter $a \in A_l$ and an infinite set $I \subseteq \mathbb{N}$ such that $W_m \prec \sigma^{k_m}(a)$ for $m \in I$. Write down each word $\sigma^n(a)$ as

$$\sigma^n(a) = L_1^{'(n)} S_1^{'(n)} \dots L_k^{'(n)} S_k^{'(n)} L_{k+1}^{'(n)} \dots L_{r_n}^{'(n)} S_{r_n}^{'(n)}$$

where $L_i^{'(n)} \in A_l^+$ and $S_i^{'(n)} \in A_s^+$. Observe also that the words $L_1^{'(n)}$ and $S_{r_n}^{'(n)}$ can be empty. Since $\sigma(S_i^{'(n)}) \in A_s^+$, the maximal length of $\{S_1^{'(n)}, \dots, S_{r_n}^{'(n)}\}$ must monotonically tend to infinity as $n \to \infty$.

Let $\tau = \sigma^{n_0 p}$, where the integers n_0 and p are defined above. Then, we decompose each word $\tau(a)$, $a \in A_l$, into blocks over short and long letters

$$\tau^n(a) = L_1^{(n)} S_1^{(n)} \dots L_k^{(n)} S_k^{(n)} L_{k+1}^{(n)} \dots L_{d_n}^{(n)} S_{d_n}^{(n)} \tag{6}$$

where $L_i^{(n)} \in A_l^+$ and $S_i^{(n)} \in A_s^+$. We notice that the maximal length of $\{S_1^{(n)}, \dots, S_{d_n}^{(n)}\}$ tends to infinity as $n \to \infty$.

Since $\tau(S_k^{(n)}) \in A_s^+$, each block $S_k^{(n)}$ is *followed by a block* $S_i^{(n+1)}$ (in symbols, $S_k^{(n)} < S_i^{(n+1)}$) in the following sense: let $a_k^{(n)}$ and $b_k^{(n)}$ denote the last and the first letters of $L_k^{(n)}$, respectively, then $S_i^{(n+1)}$ is the maximal factor of $\tau(a_k^{(n)} S_k^{(n)} b_{k+1}^{(n)}) = \tau(a_k^{(n)})\tau(S_k^{(n)})\tau(b_{k+1}^{(n)})$ over A_s^+ that contains $\tau(S_k^{(n)})$. In other words, $S_i^{(n+1)}$ is the maximal factor of $\tau^{n+1}(a)$ over A_s^+ that appears between $a_i^{(n+1)}$ and $b_{i+1}^{(n+1)}$.

(2) For N sufficiently large, there exists $1 \le k_0 \le d_N$ (see (6) for the definition of d_N) such that the block $S_{k_0}^{(N)}$ is followed by blocks $S_{k_i}^{(N+i)}$,

$$S_{k_0}^{(N)} < S_{k_1}^{(N+1)} < S_{k_2}^{(N+2)} < \dots$$

with $|S_{k_i}^{(N+i)}| \to \infty$ as $i \to \infty$.

Therefore, we see from the above argument that

$$\begin{aligned} \tau^i(a_{k_0}^{(N)}) &= \boxed{*}\, a_{k_i}^{(N+i)} V_i, \\ \tau^i(b_{k_0+1}^{(N)}) &= Z_i b_{k_i+1}^{(N+i)} \boxed{**} \end{aligned}$$

where $V_i, Z_i \in A_s^*$ and $\boxed{*}$, $\boxed{**}$ are some words. Then we can find positive integers q and t such that

$$\begin{aligned} a_{k_t}^{(N+t)} &= a_{k_{t+ql}}^{(N+t+ql)}, \\ b_{k_t+1}^{(N+t)} &= b_{k_{t+ql}+1}^{(N+t+ql)} \end{aligned}$$

for all $l \geq 0$.

(3) Since $|S_{k_i}^{(N+i)}| \to \infty$, we have that $a_{k_i}^{(N+i)}$ or $b_{k_i+1}^{(N+i)}$ adds to $S_{k_0}^{(N)}$ new blocks over A_s^+, i.e., $|V_i| \to \infty$ or $|Z_i| \to \infty$ as $i \to \infty$. For definiteness, we assume that so does $a_{k_i}^{(N+i)}$. Therefore,

$$\tau^q(a_{k_t}^{(N+t)}) = \boxed{*}a_{k_{t+q}}^{(N+t+q)}Q = \boxed{*}a_{k_t}^{(N+t)}Q$$

and, inductively,

$$\tau^{lq}(a_{k_t}^{(N+t)}) = \boxed{**}a_{k_t}^{(N+t)}Q\tau^q(Q)\dots\tau^{q(l-1)}(Q),\ l \geq 1,$$

where $Q \in A_s^+$ and $\boxed{*}$, $\boxed{**}$ are some words.

It follows from the definition of τ that $\tau^l(Q) = \tau(Q)$ for every $l \geq 1$. Therefore, X_σ contains a periodic sequence $\tau(Q)^\infty$, which is impossible. $\square$

Proposition 5.6 *Let $\sigma : A \to A^+$ be an aperiodic substitution. Then (X_σ, T_σ) has no more than $|A|$ minimal components.*

Proof. Assume the converse. Set $K = |A|$. Take any $K+1$ minimal components $Z_0, \dots, Z_K$ and let $z_i \in Z_i$, $i = 0, \dots, K$.

We claim that there exist an infinite set $I \subseteq \mathbb{N}$ and letters $b_j \in A_l$, $j = 0, \dots, K$, such that $\sigma^i(b_j) \prec z_j$ for all $i \in I$, $j = 0, \dots, K$.

Indeed, by Proposition 5.5, find $M > 0$ such that any word $w \in L(X_\sigma)$ of length at least M contains a letter from A_l. Given $n > 0$, take $m > M||\sigma^n||$. By definition of X_σ, $z_0[-m, m]$ is a factor of $\sigma^k(a)$ for some $a \in A$ and k. Since the function $n \mapsto ||\sigma^n||$ is not decreasing, we get that $k > n$. Let $\sigma^{k-n}(a) = a_0 \dots a_{d-1}$ with $a_i \in A$. We can choose the maximal interval $[i, j]$ of $[0, d-1]$ such that $\sigma^n(a_i \dots a_j)$ appears in $z_0[-m, m]$. The choice of m guarantees us that $j - i \geq M$. Therefore, at least one of the letters a_r belongs to A_l, $i \leq r \leq j$.

Thus, for all $n \in \mathbb{N}$ there is $a_n \in A_l$ such that $\sigma^n(a_n)$ appears in z_0. Therefore, there is an infinite set $I_0 \subseteq \mathbb{N}$ and a letter $b_0 \in A_l$ such that $\sigma^i(b_0)$ appears in z_0 for all $i \in I_0$. Analogously, for all $i \in I_0$ there is $a_i \in A_l$ such that $\sigma^i(a_i)$ appears in z_1. Therefore, there is an infinite set $I_2 \subseteq I_1$ and a letter $b_1 \in A_l$ such that $\sigma^i(b_1)$ appears in z_1 for all $i \in I_1$. Repeating the argument, we find $I = I_K$ and letters $b_0, \dots, b_{K-1}$. This proves the claim.

Since we have only K letters, there is a letter, say $a \in A_l$, such that $\sigma^i(a)$ appears in two distinct sequences z_l and z_d for all $i \in I$. Fix any $i \in I$. Denote by $E[i]$ the integer part of $|\sigma^i(a)|/2$. Note that $|\sigma^i(a)| \to \infty$. By shifting z_l and z_d, if necessary, we can assume that $\sigma^i(a)$ appears in z_l and z_d in such a way that the $E[i]$-th coordinate of $\sigma^i(a)$ is aligned with zero coordinates of z_l and z_d. Therefore, the d-distance between the compact sets Z_l and Z_d is less than $1/2^{E[i]}$, for any $i \in I$, which is impossible. $\square$

Remark 5.7. It immediately follows from the proof of Proposition 4.6 that the result is still true if we replace the aperiodicity of σ by the condition that $|\sigma^n| \to \infty$ as $n \to \infty$.

Definition 5.8. Let $\sigma : A \to A^+$ be a substitution, $a \in A$, $k > 0$. The words of the form $\sigma^k(a)$ will be called *k-words*. Let $n > k$ and $\sigma^{n-k}(a) = a_0 \dots a_{m-1}$ with $a_i \in A$. Then $\sigma^n(a) = \sigma^k(a_0) \dots \sigma^k(a_{m-1})$. We will say that the n-word $\sigma^n(a)$ is *naturally decomposed into k-words* $[\sigma^k(a_0), \dots, \sigma^k(a_{m-1})]$.

Remark 5.9. Suppose that an n-word $\sigma^n(a)$ is naturally decomposed into k-words $[\sigma^k(a_0), \dots, \sigma^k(a_{m-1})]$ and into $k+1$-words $[\sigma^{k+1}(c_0), \dots, \sigma^{k+1}(c_{l-1})]$, i.e.,

$$\sigma^n(a) = \sigma^k(\sigma(c_0)) \dots \sigma^k(\sigma(c_{l-1})) = \sigma^k(a_0) \dots \sigma^k(a_{m-1}).$$

Since the decompositions are natural, we get that each letter a_i appears as a factor of some $\sigma(c_j)$.

The following result shows that each element of X_σ can be written as a concatenation of 1-words.

Proposition 5.10 *Let $\sigma : A \to A^+$ be a substitution. Then for every $x \in X_\sigma$ there exist $y \in X_\sigma$ and $i \in \{0, \dots, |\sigma(y[0])| - 1\}$ such that $x = T_\sigma^i \sigma(y)$.*

Proof. *I.* Take any $x \in X_\sigma$. For each $n \geq 1$, find $m_n > n||\sigma^n||$. By definition of X_σ, there are $v^{(n)} \in A$ and $k_n > 0$ such that $x[-m_n, m_n]$ is a factor of $\sigma^{k_n}(v^{(n)})$. It is evident that $k_n > n$. Let $\sigma^{k_n - n}(v^{(n)}) = v_1^{(n)} \dots v_{d_n}^{(n)}$ with $v_i^{(n)} \in A$. Hence, the word $x[-m_n, m_n]$ is a factor of the word $\sigma^n(v_1^{(n)}) \dots \sigma^n(v_{d_n}^{(n)}) = \sigma^{k_n}(v^{(n)})$ which is considered as the concatenation of n-words. Take the maximal interval $[i_n, j_n] \subseteq \{1, \dots, d_n\}$ so that the word $\sigma^n(v_{i_n}^{(n)})\sigma^n(v_{i_n+1}^{(n)}) \dots \sigma^n(v_{j_n}^{(n)})$ appears as a factor of $x[-m_n, m_n]$. That is

$$x[-l_n, |\sigma^n(v_{i_n}^{(n)} \dots v_{j_n}^{(n)})| - 1 - l_n] = \sigma^n(v_{i_n}^{(n)} \dots v_{j_n}^{(n)}),$$

where l_n is defined as the natural position of occurrence of $x[0]$ within the word $\sigma^n(v_{i_n}^{(n)}) \dots \sigma^n(v_{j_n}^{(n)})$. Setting $w_n = v_{i_n}^{(n)} \dots v_{j_n}^{(n)}$, we get that

$$x[-l_n, |\sigma^n(w_n)| - 1 - l_n] = \sigma^n(w_n) \text{ for all } n \geq 1 \tag{7}$$

and $l_n \to \infty$, $(|\sigma^n(w_n)| - l_n) \to \infty$ as $n \to \infty$.

II. For each n and $i = i_n, \dots, j_n$, take the natural decomposition of the n-word $\sigma^n(v_i^{(n)})$ into 1-words and write down all of these 1-words from left to right, say

$$\sigma^n(w_n) = \sigma^n(v_{i_n}^{(n)}) \dots \sigma^n(v_{j_n}^{(n)}) = \sigma(y_{-s_n}^{(n)}) \dots \sigma(y_{d_n}^{(n)}).$$

Enumerate these 1-words by the following rule: as all of them naturally appear in the sequence x, we set $\sigma(y_0^{(n)})$ to be the 1-word that naturally covers $x[0]$; $\sigma(y_{-s}^{(n)})$ is the s-th 1-word on the left of $\sigma(y_0^{(n)})$; and $\sigma(y_s^{(n)})$ is the s-th 1-word on the right of $\sigma(y_0^{(n)})$ if $s > 0$.

For every $n \geq 1$, define the integer $f_0^{(n)} \leq 0$ as the position of the natural occurrence of 1-word $\sigma(y_0^{(n)})$ within x, i.e.,

$$x[f_0^{(n)}, |\sigma(y_0^{(n)})| - 1 + f_0^{(n)}] = \sigma(y_0^{(n)}).$$

III. Considering 1-words $\sigma(y_0^{(n)})$ and the positions of their occurrences in x, we find an infinite subset $I_0 \subseteq \mathbb{N}$ such that

$$y_0^{(n)} = y_0^{(n')} \text{ and } f_0^{(n)} = f_0^{(n')} \text{ for any } n, n' \in I_0.$$

Then by induction on k, we find an infinite set $I_k \subseteq I_{k-1} \subseteq \ldots \subseteq I_0$ such that

$$y_i^{(n)} = y_i^{(n')} \text{ for any } n, n' \in I_k \text{ and } i = -k, \ldots, k.$$

This defines a two-sided sequence

$$y = \{\ldots, y_{-k}^{(n_k)}, y_{-k+1}^{(n_{k-1})}, \ldots, y_0^{(n_0)}, y_1^{(n_1)}, \ldots, y_k^{(n_k)}, \ldots\} \text{ with } n_k \in I_k.$$

Observe that $y \in X_\sigma$ and $x = T_\sigma^i \sigma(y)$, where $i = -f_0^{(n_0)}$. □

<table>
<tr><td>...</td><td>x_{-3}</td><td>x_{-2}</td><td>x_{-1}</td><td>x_0</td><td>x_1</td><td>x_2</td><td>x_3</td><td>x_4</td><td>...</td></tr>
<tr><td>...</td><td colspan="2">$\sigma(y_{-1})$</td><td colspan="3">$\sigma(y_0)$</td><td colspan="3">$\sigma(y_1)$</td><td>...</td></tr>
</table>

Decomposition into 1-words

Remark 5.11. As an immediate corollary of Proposition 5.10 we get that for any $n > 0$ and any $x \in X_\sigma$ there exists $y \in X_\sigma$ and $i \in \{0, \ldots, |\sigma^n(y[0])| - 1\}$ such that $x = T_\sigma^i \sigma^n(y)$.

Definition 5.12. We say that a substitution is *recognizable* if for each $x \in X_\sigma$ there exist a unique $y \in X_\sigma$ and unique $i \in \{0, \ldots, |\sigma(y[0])| - 1\}$ such that

$$x = T_\sigma^i \sigma(y). \tag{8}$$

Remark 5.13. We note that in the theory of primitive substitutions it was a long-standing problem to establish the uniqueness of this representation. It follows from the works [Hos] or [Que] that under the assumption of *bilateral recognizability* (see [Mos1]) and injectivity of the substitution on the alphabet, each aperiodic primitive substitution is recognizable in our sense. However, due to the works of Mossé [Mos1] and [Mos2] it became clear that, in fact, each aperiodic primitive substitution is recognizable.

Now we will show that an arbitrary aperiodic substitution is recognizable. Our proof involves the usage of Downarowicz-Maass' techniques developed in Section 3. So, first of all, we introduce agreeable families of j-symbols.

Definition 5.14. Let $\sigma : A \to A^+$ be a substitution. Set $A_{i-1} = A$ and $\sigma_i = \sigma$, for all $i \geq 1$. Denote by $\mathcal{A}_j$ the family of j-symbols determined by the alphabets $\{A_i\}_{i \geq 0}$ and the maps $\{\sigma_i\}_{i \geq 1}$ as in Definition 3.2.

To make the explanation more comprehensible, we will label the boxes from a row i of j-symbol $[a]_j \in \mathcal{A}_j$ by the symbols $\sigma^i(b)$, $b \in A$, instead of just $b \in A$.

<table>
<tr><td>b_0</td><td>b_1</td><td>b_2</td><td>b_3</td><td>$\ldots$</td><td>b_{m-3}</td><td>b_{m-2}</td><td>b_{m-1}</td></tr>
<tr><td colspan="2">$\sigma(c_0)$</td><td colspan="2">$\sigma(c_1)$</td><td>$\ldots$</td><td colspan="3">$\sigma(c_{d-1})$</td></tr>
<tr><td colspan="8">$\vdots$</td></tr>
<tr><td colspan="8">$\sigma^j(a)$</td></tr>
</table>

j-symbol

Denote by $\mathcal{Z}_j$ the set of all j-sequences, see Definition 3.7. Let $T : \mathcal{Z}_j \to \mathcal{Z}_j$ again denote the shift.

Definition 5.15. For $y \in X_\sigma$ and $k \in \mathbb{Z}$, denote by $[y]_j^k = \{[y(i)]_j\}_{i \in \mathbb{Z}}$ the j-sequence obtained by concatenation of j-symbols $[y(i)]_j$, $i \in \mathbb{Z}$, where the j-symbol $[y(0)]_j$ appears in $[y]_j^k$ at the position k. In other words, $[y]_j^k$ is a matrix whose j-th row is a concatenation of boxes labeled by $\sigma^j(y(i))$, $i \in \mathbb{Z}$, from left to right such that the box labeled by $\sigma^j(y(0))$ starts at the column k.

Denote by Ω_j the closed shift-invariant subset of $\mathcal{Z}_j$ generated by all j-sequences $[y]_j^k$ with $y \in X_\sigma$ and $k \in \mathbb{Z}$.

The proof of the following lemma is analogous to that of Proposition 5.6, so we omit it.

Lemma 5.16 *(Ω_j, T) has no more than $|A|$ minimal components for any $j \geq 0$.*

Now we are ready to show that each substitutional dynamical system without periodic points is recognizable. Note that the proof uses the ideas of that of Theorem 4.8.

Theorem 5.17 *Each aperiodic substitution $\sigma : A \to A^+$ is recognizable.*

Proof. Assume that σ is not recognizable. Then we have a situation which is shown on the picture below, i.e., there exist $y_{-1}, x_{-1} \in X_\sigma$, $j_y \in \{0, \ldots, |\sigma(y_{-1}(0))| - 1\}$, and $j_x \in \{0, \ldots, |\sigma(x_{-1}(0))| - 1\}$ with $y_0 = T^{j_y}\sigma(y_{-1}) = T^{j_x}\sigma(x_{-1}) = x_0$ and $j_y \neq j_x$ or $y_{-1} \neq x_{-1}$.

<table>
<tr><td>$\ldots$</td><td>$x(-3)$</td><td>$x(-2)$</td><td>$x(-1)$</td><td>$x(0)$</td><td>$x(1)$</td><td>$x(2)$</td><td>$x(3)$</td><td>$x(4)$</td><td>$x(5)$</td><td>$\ldots$</td></tr>
<tr><td colspan="2">$\ldots$</td><td colspan="2">$\sigma(y_{-1}(-1))$</td><td colspan="4">$\sigma(y_{-1}(0))$</td><td colspan="2">$\sigma(y_{-1}(1))$</td><td>$\ldots$</td></tr>
<tr><td>$\ldots$</td><td colspan="2">$\sigma(x_{-1}(-1))$</td><td colspan="2">$\sigma(x_{-1}(0))$</td><td colspan="4">$\sigma(x_{-1}(1))$</td><td colspan="2">$\ldots$</td></tr>
</table>

(1) Assume that $y_{-1} = x_{-1}$. Hence $j_y \neq j_x$. Thus, $T^{j_x}\sigma(y_{-1}) = T^{j_y}\sigma(y_{-1})$. This shows that $\sigma(y_{-1})$ is a periodic point, which is impossible.

(2) Thus, $y_{-1} \neq x_{-1}$. By Proposition 5.10, we can find for each $i \geq 1$ some $x_{-i}, y_{-i} \in X_\sigma$ such that $T^{k_i}\sigma(x_{-i}) = x_{-i+1}$ and $T^{l_i}\sigma(y_{-i}) = y_{-i+1}$ for some $l_i \in \{0, \ldots, |\sigma(y_{-i}(0))| - 1\}$ and $k_i \in \{0, \ldots, |\sigma(x_{-i}(0))| - 1\}$. Observe that $x_{-i} \neq y_{-i}$, $i \geq 1$. Set also $y_i = x_i = \sigma^i(x_0)$, $i \geq 1$.

(3) Set $K = |A|$ and $L = K^2 + K$. It follows from Proposition 3.10 that there are integers

$$0 \leq i_0 < j_0 \leq i_1 < j_1 \leq \ldots \leq i_L < j_L = M$$

such that any pair of M-sequences from Ω_M with depth i_l has no common j_l-cuts and, therefore, no common M-cuts, $l = 0, \ldots, L$.

For each $i = 0, \ldots, M$, define M-sequences

$$X_i = [x_{i-M}]_M^{n_i} \text{ and } Y_i = [y_{i-M}]_M^{m_i}$$

(see Definition 5.15) where n_i and m_i are unique integers such that

$$T_\sigma^{n_i}\sigma^M(x_{i-M}) = x_i = y_i = T_\sigma^{m_i}\sigma^M(y_{i-M}).$$

In other words, the M-sequence X_i is built by the rule: the row 0 consists of concatenated boxes $x_i(j)$, $j \in \mathbb{Z}$; the row 1 consists of concatenated boxes $\sigma(x_{i-1}(j))$ aligned in such a way that if we decompose the row 1 into the letters we get the equality between rows 0 and 1; and so on; the bottom line is the concatenation of boxes $\{\sigma^M(x_{i-M}[j])\}$, $j \in \mathbb{Z}$, such that whenever we decompose the line M into the letters we get an equality between rows 0 and M. Note that the M-symbol $\sigma^M(x_{i-M}[0])$ appears in X_i at the position $-n_i$. The M-sequence Y_i is built up by the same rule.

(4) It is not hard to see that for each $l = 0, \ldots, L$ the pair (X_{i_l}, Y_{i_l}) has the depth i_l and, therefore, has no common j_l-cuts.

Let $E = E(\Omega_M, T)$ be the enveloping semigroup of (Ω_M, T), see Definition 4.7. By definition, $Ex = \{\gamma(x) : \gamma \in E\} = \overline{Orb_T(x)}$. It follows that Ex contains at least one minimal T-component. It follows from Lemma 5.16 that there is at most K minimal T-components, say $Z_0, \ldots, Z_{K-1}$. Fix any M-sequence $Q_i \in Z_i$, $i = 0, \ldots, K - 1$.

For each pair (X_{i_l}, Y_{i_l}), $l = 0, \ldots, L$, find an element $\gamma_{i_l} \in E$ such that $\gamma(X_{i_l}) = Q_j$ for some Q_j, $j = j(i_l)$.

(4-a) Since γ_{i_l} is the pointwise limit of a sequence (T^{n_k}), the sequences $\gamma_{i_l}(X_{i_l})$ and $\gamma_{i_l}(Y_{i_l})$ remain i_l-compatible. The fact that the pair (X_{i_l}, Y_{i_l}) has no common j_l-cuts implies that $(\gamma_{i_l}(X_{i_l}), \gamma_{i_l}(Y_{i_l}))$ have no common j_l-cuts as well. For otherwise, $T^{n_k}(X_{i_l})$ and $T^{n_k}(Y_{i_l})$ have a common j_l-cut for all k big enough, which is impossible.

(4-b) Consider the pairs $(\gamma_{i_l}(X_{i_l}), \gamma_{i_l}(Y_{i_l}))$, $l \in [0, K^2 + K]$. As every $\gamma_{i_l}(X_{i_l})$ coincides with some of $\{Q_0, \ldots, Q_{K-1}\}$, we can choose K^2 pairs $(\gamma_i(X_{i_l}), \gamma_i(Y_{i_l}))$, $l \in I$, such that $\gamma_{i_l}(X_{i_l})$ are the same for all $l \in I$, $|I| = K^2$, say $\gamma_{i_l}(X_{i_l}) = Q_{k_0}$ when $l \in I$.

Since each pair $(\gamma_{i_l}(X_{i_l}), \gamma_{i_l}(Y_{i_l}))$ is i_l-compatible and has no common j_l-cuts, the choice of the set I and integers $0 \leq i_0 < j_0 \leq \ldots\ldots i_L < j_L$ guarantees us that each pair $(\gamma_{i_l}(Y_{i_l}), \gamma_{i_{l'}}(Y_{i_{l'}}))$, $l, l' \in I$ and $l < l'$, is 0-compatible and has no common $j_{l'}$-cuts.

(4-c) Setting $Q'_l = \gamma_{i_l}(Y_{i_l})$ for $l \in I$, we obtain a family $\{Q'_l : l \in I\} \cup \{Q_{k_0}\}$ of $K^2 + 1$ M-sequences which are 0-compatible and have no common M-cuts. Applying Proposition 3.12, we get that $\pi_0(Q_{k_0})$ is eventually periodic. It follows from the minimality of Q_{k_0} that $\pi_0(Q_{k_0}) \in X_\sigma$ is periodic, which is a contradiction. □

For each $a \in A$, set $[a] = \{x \in X_\sigma : x_0 = a\}$. Note that, in general, the set $[a]$ could be empty. However, the following result is true even in the case when some of the sets $[a]$, $a \in A$, are empty.

Corollary 5.18 *Let $\sigma : A \to A^+$ be an aperiodic substitution. Then for every $n \geq 0$*

$$\mathcal{P}_n = \{T^i_\sigma \sigma^n([a]) \ : \ a \in A \text{ and } 0 \leq i < |\sigma^n(a)|\}$$

is a clopen partition of X_σ. Furthermore, the sequence of partitions $\{\mathcal{P}_n\}$ is nested.

Proof. First of all, observe that $\sigma^n : X_\sigma \to X_\sigma$ is a continuous map. It follows from Proposition 5.10 that the closed set $\sigma^n(X_\sigma)$ meets each T_σ-orbit and consists of recurrent points. Note that the return time of each $\sigma^n(y)$, $y \in X_\sigma$ to $\sigma^n(X_\sigma)$ is at most $|\sigma^n(y[0])|$. Theorem 5.17 implies that the map $\sigma^n : X_\sigma \to X_\sigma$ is one-to-one and the first return time of each $\sigma^n(y) \in \sigma^n(X_\sigma)$ to $\sigma^n(X_\sigma)$ is exactly $|\sigma^n(y[0])|$. This shows that $\mathcal{P}_n$ is a finite partition of X_σ into closed sets and, therefore, into clopen sets. The fact that the sequence of partitions $\{\mathcal{P}_n\}$ is nested is proved in [DHS, Proposition 14]. □

Remark 5.19. Observe that, in general, the sequence of K-R partitions $\{\mathcal{P}_n\}$ may not generate the topology of X_σ.

6 Stationary Bratteli-Vershik systems vs. aperiodic substitutions

In the section we show that the class of aperiodic substitutional systems coincide with the class of expansive Vershik maps of stationary ordered Bratteli diagrams.

We recall that Bratteli-Vershik models of substitutional dynamical systems were constructed for primitive substitutions in the papers [For] and [DHS]. We also refer the reader to the papers [CE1], [CE2], [Yua1], and [Yua2], where related topics such as various dimension groups and invariant measures for substitutional dynamical systems are considered. It is worthwhile to mention the pioneering paper by Ferenczi [Fer] where the study of substitutions on infinite alphabets was initiated. We observe that these systems can be thought as aperiodic homeomorphisms of zero-dimensional Polish spaces.

6.1 From Bratteli diagrams to substitutional systems.

We start this subsection with the definition of a stationary Bratteli diagram.

Definition 6.1. (1) A Bratteli diagram $B = (V, E)$ is *stationary* if $k = |V_1| = |V_2| = \dots$ and if (by an appropriate labeling of the vertices) the incidence matrix between levels n and $n+1$ is the same $k \times k$ matrix C for all $n = 1, 2, \dots$. In other words, beyond level 1 the diagram repeats. Clearly, we can label the vertices in V_n as $V_n(a_1), \dots, V_n(a_k)$, where $A = \{a_1, \dots, a_k\}$ is a set of k distinct symbols.

(2) A Bratteli diagram $B = (V, E, \leq)$ is *stationary ordered* if (V, E) is stationary and the ordering '$\leq$' on the edges with range $V_n(a_i)$ does not depend on $n \geq 2$, for all $i = 1, \dots, k$.

(3) Let $B = (V, E, \leq)$ be a stationary ordered Bratteli diagram and V_n denote the set of vertices at level n, $n \geq 0$. Choose a stationary labeling of V_n by an alphabet A, i.e. $V_n = \{V_n(a) \mid a \in A\}$ for $n > 0$. For every letter $a \in A$, consider the ordered set $(e_1, \dots, e_k)$ of edges that range at $V_n(a)$, $n \geq 2$, and let $(a_1, \dots, a_k)$ be the ordered set of the labels of the sources of these edges with respect to the ordering '$\leq$'. The map $a \mapsto a_1 \dots a_k$ from A to A^+ does not depend on n and therefore determines a substitution called the *substitution read on* B.

The following result shows that stationary diagrams can have only a finite number of minimal and maximal paths. See also Propositions 4.6 and 5.6 for similar results.

Proposition 6.2 *Let $B = (V, E, \leq)$ be a stationary ordered Bratteli diagram. Then B has a finite number of minimal and maximal paths.*

Proof. Let σ be the substitution read on B. Set

$$A_0 = \{a \in A \ : \ \text{there is } n > 0 \ (\sigma^n(a) \text{ begins with } a)\}$$

and

$$A_1 = \{b \in A \ : \ \text{there is } n > 0 \ (\sigma^n(b) \text{ ends with } b)\}.$$

For each $a \in A_0$, let n_a be an integer such that $\sigma^{n_a}(a)$ begins with a. Analogously, for each $b \in A_1$, let n_b be an integer such that $\sigma^{n_b}(b)$ ends with b.

Setting $p = \prod_{a \in A_0} n_a \times \prod_{b \in A_1} n_b$, we see that for any $c \in A$ the word $\sigma^n(c)$ begins (ends) with c for some $n > 0$ if and only if $\sigma^p(c)$ begins (ends) with c.

Now consider a minimal path x of B. Since we have exactly $|A|$ vertices at each level, we can find a letter $a \in A$ and an infinite set $I \subseteq \mathbb{N}$ such that x goes through the vertex a at levels k with $k \in I$. In particular, this means that $\sigma^p(a)$ begins with a. We can write down each $k \in I$ as

$$k = pl_k + m_k \text{ with } l_k \geq 0 \text{ and } 0 \leq m_k < p.$$

Find an infinite set $J \subseteq I$ such that $m_k = m_{k'}$ for $k, k' \in J$. It follows that x goes through the vertex labeled by a at levels $m + np$, $n \geq 1$, where $m = m_k$

for some $k \in J$. This, in particular, shows that there is only a finite number of minimal paths. The proof of the result for maximal paths is analogous. $\square$

The following proposition shows that the expansiveness of Vershik maps defined on stationary Bratteli diagrams is already seen at the first level.

Proposition 6.3 *Suppose that $B = (V, E, \leq)$ is a stationary Bratteli diagram with continuous Vershik map. If x and y from X_B are 1-compatible, then for any $i \geq 1$ there exists a pair of i-compatible elements. In particular, this means that such a Vershik map is not expansive.*

Proof. We consider a map $f : X_B \to X_B$, which was originally defined in [DHS]. For $x \in X_B$, let x_n be the label of the edge between levels $n-1$ and n the path x goes through. For all $n \geq 3$, let y_n be x_{n-1} and (y_1, y_2) be the minimal path connecting v_0 and $s(y_3)$. Set $f(x) = y$, where $y = (y_1, y_2, \ldots)$.

Now it is not hard to see that if z_1 and z_2 are 1-compatible, then $f^i(z_1)$ and $f^i(z_2)$ are i-compatible. $\square$

The following theorem generalizes the main result of [DHS] to any aperiodic Vershik map.

Theorem 6.4 *Suppose that (X_B, φ_B) is an aperiodic Bratteli-Vershik system with B a stationary ordered Bratteli diagram and X_B is perfect. Then the system (X_B, φ_B) is homeomorphic to an aperiodic substitutional dynamical system if and only if no restriction of φ_B to a minimal component is isomorphic to an odometer.*

Proof. *(I)* Assume first that the diagram B has only single edges between the top vertex and the vertices of the first level. Let A be the labeling of vertices of B as in Definition 6.1 and $\sigma : A \to A^+$ be the substitution read on B. Define a map $\pi : X_B \to A^{\mathbb{Z}}$ as follows

$$\pi(x)_k = a \text{ if and only if } \varphi_B^k(x) \text{ passes through } V_1(a).$$

Observe that $\pi = \pi_1$, where π_1 maps each $x \in X_B$ to the row 1 of its matrix, see Section 4. Clearly, π is continuous and

$$\pi \circ \varphi_B = T \circ \pi, \tag{9}$$

where T is the shift on $A^{\mathbb{Z}}$.

(1) We claim that $\pi(X_B) \subseteq X_\sigma$. Indeed, to check this, it is sufficient to prove that for every $x \in X_B$ and any n one has $\pi(x)[-n, n] \in L(\sigma)$.

Assume that $x \in X_B$ is cofinal neither to a maximal nor to a minimal path. Let $[x]$ denote the matrix that is obtained by concatenation of j-symbols determined by x, see Section 4. Then there is $j > 0$ such that the j-symbol, say v, from $[x]$ crossing 0 column 'covers' coordinates $[-n, n]$ of the first line. This implies that $\pi(x)[-n, n]$ is a factor of $\sigma^j(v)$.

By Proposition 6.2 the diagram has only a finite number of maximal and minimal paths. Since X_B is perfect, the set $X_B \setminus Orb_{\varphi_B}(X_{\max} \cup X_{\min})$ is dense in X_B. It follows from the continuity of π that $\pi(X_B) \subseteq X_\sigma$.

(2) We assert that $\pi(X_B)$ is dense in X_σ. Indeed, consider the cylinder set $C = \{x \in X_\sigma : x[n, n + |\sigma^j(a)| - 1] = \sigma^j(a)\}$ with $a \in A$, $j \geq 1$, and $n \in \mathbb{Z}$. By Proposition 5.10, such cylinder sets generate the topology on X_σ. Thus, we need to show that there is $x \in X_B$ such that $\pi(x) \in C$. Observe that by (9), it is sufficient to find $x \in X_B$ and $k \in \mathbb{Z}$ such that $\pi(x)[k, k + |\sigma^j(a)| - 1] = \sigma^j(a)$.

Let $(y_1, \ldots, y_j)$ stand for the minimal path connecting v_0 to $V_j(a)$. If $x \in U(y_1, \ldots, y_j)$, then the j-symbol $[a]_j$ appears in the matrix $[x]$ at the coordinate 0. Therefore, $\pi(x)[0, |\sigma^j(a)| - 1] = \sigma^j(a)$.

(3) It follows from the continuity of π that $\pi(X_B) = X_\sigma$. Thus, (X_σ, T_σ) is a factor of (X_B, φ_B).

(4) Suppose that the restriction of φ_B onto one of its minimal components is homeomorphic to an odometer. Then φ_B is obviously non-expansive[2]. Due to the fact that each substitutional dynamical system is expansive, the homeomorphism φ_B cannot be homeomorphic to a substitutional systems.

Conversely, if none of the restrictions of φ_B to minimal components is homeomorphic to an odometer, then by Theorem 4.8 (X_B, φ_B) is expansive. By Proposition 6.3 the map π is injective. This shows that (X_B, φ_B) and (X_σ, T_σ) are homeomorphic.

(II) Now, let B be an arbitrary ordered Bratteli diagram with the expansive aperiodic Vershik map $\varphi_B : X_B \to X_B$. There are two ways of realization of (X_B, φ_B) as a substitutional dynamical system. The first one is to build a diagram equivalent to B that has only simple edges on the first level, and then apply part (I). The second one is to construct the substitutional dynamical system directly from the diagram B. We will exploit the second approach which was first applied in [DHS, Proposition 23] for primitive substitutions.

(1) Denote by A the labeling of vertices from V_n $(n \geq 1)$ as in Definition 6.1. For each vertex $a \in A$, let n_a be the number of edges between the vertex $V_1(a)$ of the first level labeled by a and the top vertex v_0.

Denote by D an ordered Bratteli diagram that coincides with B everywhere, but has only one edge between each vertex $V_1(a)$ and the top vertex v_0. Let φ_D be the restriction of the Vershik map φ_B to X_D considered as a clopen subset of X_B. Note that (X_D, φ_D) is an induced system of (X_B, φ_B). It was proved in (I) that (X_D, φ_D) is homeomorphic to the substitutional dynamical system (X_σ, T_σ), where σ is the substitution read on D.

(2) Set $m_a = |\sigma(a)|$, $a \in A$. Since the Vershik map φ_D is aperiodic, we obtain that $|\sigma^n| \to \infty$ where the substitution σ is read on the diagram D. Thus, we can assume, substituting a power of σ for σ if needed, that $m_a \geq n_a$ for every $a \in A$.

Let (X_1, T) denote the factor of (X_B, φ_B) obtained by 1-sequence coding of elements of X_B, see Section 4. Let

$$B_1 = \{[a]_1(i) : a \in A,\ i = 0, \ldots, n_a - 1\}.$$

---

[2] It can be deduced, in particular, from Theorem 5.23 of [Wal] and the existence of Kakutani-Rokhlin partitions for the system that generate the topology.

One can see that (X_1, T) is a subshift over the alphabet B_1. By Proposition 6.3, the factor map $\pi_1 : X_B \to X_1$ is injective, so the systems (X_1, T) and (X_B, φ_B) are homeomorphic. Consider the clopen set of X_1

$$U = \bigcup_{a \in A} [a](0).$$

Clearly, the return time of any point from $[a]_1(0)$ to U is n_a. We observe that (X_σ, T_σ) is homeomorphic to $(X_1|_U, T_U)$ and the homeomorphism is implemented by the map $\xi : A \to B_1^+$ where $\xi(a) = [a]_1(0) \dots [a]_1(n_a - 1)$.

(3) Define the alphabet $E := \{(a, i) : a \in A,\ 0 \le i \le n_a - 1\}$ and the map $\psi : A \to E^+$ by

$$\psi(a) = (a, 0)(a, 1) \dots (a, n_a - 1).$$

Let $\tau : E \to E^+$ be a substitution given by

$$\tau(a, i) = \begin{cases} \psi(\sigma(a)_i) & \text{if } 1 \le i < n_a - 1 \\ \psi(\sigma(a)_{[n_a - 1, m_a)}) & \text{if } i = n_a - 1. \end{cases}$$

Take any $a \in A$, then $\tau(\psi(a)) = \tau(a, 0) \dots \tau(a, n_a - 1) = \psi(\sigma(a))$. Therefore,

$$\tau^n \circ \psi = \psi \circ \sigma^n \text{ for any } n \ge 0. \tag{10}$$

For the substitution σ read on the diagram and every $a \in A$ there are $b \in A$ such that a occurs in $\sigma(b)$, say, at position i. Therefore, $\psi(a) = \psi(\sigma(b)_i)$ is a factor of $\tau((b, i))$. Thus, $\psi(a) \in L(\tau)$ for every $a \in A$. By (10), we have that $\psi(L(\sigma)) \subseteq L(\tau)$. This shows that $\psi(X_\sigma) \subset X_\tau$.

Consider the clopen subset Q of X_τ given by

$$Q = \bigsqcup_{a \in A} [\psi(a)].$$

Clearly, $\psi(X_\sigma) \subseteq Q$.

It follows from Proposition 5.10 that for every $x \in X_\tau$ there are $z \in X_\tau$ and $0 \le k < |\tau(z[0])|$ such that $x = T_\tau^k \tau(z)$. Therefore, by definition of τ, for every $x \in X_\tau$ there exist $y \in A^{\mathbb{Z}}$ and $0 \le k < |\psi(y[0])|$ such that $x = T_\tau^k \psi(y)$. The definition of ψ implies that such y and k are unique.

Consider any $x \in Q$. Take the unique $y \in A^{\mathbb{Z}}$ such that $\psi(y) = x$. It follows from the definition of X_τ that every $\psi(y[-k, k])$ is a factor of $\tau^n((a, i))$ for some $(a, i) \in E$. Let $a = \sigma(b)_j$ for some $b \in A$ and $j = 0, \dots, m_b - 1$, i.e., $(a, i) \prec \psi(\sigma(b))$. It follows that $\tau^n((a, i))$ is a factor of $\tau^n \circ \psi(\sigma(b)) = \psi(\sigma^{n+1}(b))$. Therefore,

$$\psi(y[-k, k]) \prec \psi(\sigma^{n+1}(b)).$$

The definition of ψ implies that if $v \not\prec w$ where $v, w \in A^+$, then $\psi(v) \not\prec \psi(w)$. Hence $y[-k, k]$ is a factor of $\sigma^{n+1}(b)$. Therefore, $y \in X_\sigma$ and $\psi(X_\sigma) = Q$. This shows that ψ implements an homeomorphism between (X_σ, T_σ) and $(X_\tau|_Q, (T_\tau)_Q)$. The following diagram illustrates the relation between the systems. The downward arrows shows that the underlying system is an induced system of the overlying one.

$$\begin{tikzcd}
 & & (X_1,T) \arrow[d] & \overset{\pi_1}{\cong} & (X_B,\varphi_B) \arrow[d] & \cong & (X_\tau,T_\tau) \arrow[d] \\
(X_D,\varphi_D) & \cong & (X_1|_U,T_U) & \overset{\xi}{\cong} & (X_\sigma,T_\sigma) & \overset{\psi}{\cong} & (X_\tau|_Q,(T_\tau)_Q)
\end{tikzcd}$$

Observe that the return time of all points from $[\psi(a)]$, $a \in A$, to Q under the action of T_τ is n_a. Thus, the map $\rho : E \to B_1$ given by $\rho((a,i)) = [a]_1(i)$ implements an homeomorphism between (X_τ, T_τ) and $(X_1,T) \cong (X_B,\varphi_B)$. $\square$

6.2 From substitutional dynamical systems to Bratteli diagrams.

In this subsection we show how one can construct a Bratteli-Vershik model for aperiodic substitutional dynamical systems which satisfy the *nesting property* (see Definition 6.5). We present a technique applicable for a wide class of substitutions including those with $|\sigma^n| \to \infty$ and various Chacon-like substitutions.

Let $\sigma : A \to A^+$ be a substitution. Denote by A_l the set of all letters $a \in A$ such that $|\sigma^n(a)| \to \infty$ as $n \to \infty$. Set also $A_s = A \setminus A_l$.

Definition 6.5. We say that a substitution $\sigma : A \to A^+$ has a *nesting*[3] *property* if either (1) for every $a \in A_l$ the word $\sigma(a)$ starts with a letter from A_l or (2) for every $a \in A_l$ the word $\sigma(a)$ ends with a letter from A_l.

Theorem 6.6 *Let $\sigma : A \to A^+$ be an aperiodic substitution with nesting property. Then the substitutional dynamical system (X_σ, T_σ) is homeomorphic to the Vershik map of a stationary Bratteli diagram.*

Proof. For an alphabet B, let B^* denote the set of all words over B including the empty word.

(1) By Proposition 5.5, find $M > 0$ such that any word $W \in L(X_\sigma)$ with $|W| \geq M$ contains a letter from A_l. Set

$$\mathcal{V} = \{W \in L(X_\sigma)\ :\ w = v_1S_1v_2S_2v_3,\ v_1,v_2,v_3 \in A_l,\ \text{and}\ S_1,S_2 \in A_s^*\}.$$

Since no block $S \in A_s^+ \cap L(X_\sigma)$ can have the length greater than M, the set $\mathcal{V}$ is finite.

Since the substitution σ has the nesting property, we have that either (i) all words $\sigma(a)$ start with letters from A_l for every $a \in A_l$, or (ii) all words $\sigma(a)$ end with letters from A_l for every $a \in A_l$.

In the first case we define the sequence of K-R partitions as follows

$$\Xi_n = \{T_\sigma^i \sigma^n([v_1S_1.v_2S_2v_3])\ :\ v_1S_1v_2S_2v_3 \in \mathcal{V},\ 0 \leq i < |\sigma^n(v_2S_2)|\},\ n \geq 0.$$

---

[3] We use the term 'nesting' as a natural sequence of K-R partitions for these substitutional systems is nested, see the proof of Theorem 6.6.

In the second case we set

$$\Xi'_n = \{T^i_\sigma \sigma^n([v_1.S_1v_2S_2v_3]) \ : \ v_1S_1v_2S_2v_3 \in \mathcal{V},\ 0 \le i < |\sigma^n(S_1v_2)|\},\ n \ge 0.$$

We will consider in the proof of the theorem the first case only. The other case is proved analogously.

(2) We claim that $\{\Xi_n\}$ is a nested sequence of clopen K-R partitions. Indeed, assume that

$$T^i_\sigma\sigma^n([v_0v_1\dots v_s|v_{s+1}\dots v_{q-1}]) \cap T^j_\sigma\sigma^n([w_0\dots w_k|w_{k+1}\dots w_{r-1}]) \neq \emptyset,$$

where $0 \le i < |\sigma^n(v_{s+1}\dots v_{q-2})|$, $0 \le j < |\sigma^n(w_{k+1}\dots w_{r-2})|$, the letters $v_0, v_{s+1}, v_{q-1}, w_0, w_{k+1}, w_{r-1}$ belong to A_l, and the remaining letters v_t, w_t are taken from A_s.

Then there exist $y \in [v_0\dots v_s|v_{s+1}\dots v_{q-1}]$ and $z \in [w_0\dots w_k|w_{k+1}\dots w_{r-1}]$ with $T^i\sigma^n(y) = T^j\sigma^n(z)$. By Theorem 5.17 and the definition of $\mathcal{V}$, we get that $y = z$ and $i = j$. This implies that $q = r$, $k = s$, and $v_t = w_t$ for $t = 0,\dots,q-1$. Thus, Ξ_n is a K-R partition of X_σ. Notice that, by continuity of σ^n, all the sets $T^i_\sigma\sigma^n([v_1S_1.v_2S_2v_3])$, $v_1S_1v_2S_2v_3 \in \mathcal{V}$, are closed. Therefore, they are clopen.

Applying σ to any $W = v_1S_1|v_2S_2v_3 \in \mathcal{V}$, we get a representation

$$\sigma(W) = w^{(1)}_1F^{(1)}_1\dots w^{(1)}_{k_1}F^{(1)}_{k_1}|w^{(2)}_1F^{(2)}_1\dots w^{(2)}_{k_2}F^{(2)}_{k_2}w^{(3)}_1F^{(3)}_1\dots w^{(3)}_{k_3}F^{(3)}_{k_3}$$

with $w^{(\cdot)}_i \in A_l$, and $F^{(\cdot)}_i \in A^*_s$. Here $\sigma(v_iS_i) = w^{(i)}_1F^{(i)}_1\dots w^{(i)}_{k_i}F^{(i)}_{k_i}$ where $i = 1, 2, 3$ and $S_3 = \emptyset$. Since σ has a nesting property, we obtain that $w^{(i)}_1 \neq \emptyset$. This representation shows that the T_σ-tower from Ξ_n with base $\sigma^n([v_1S_1.v_2S_2v_3])$ consequently intersects the T_σ-towers of Ξ_{n-1} with bases:

$$\sigma^{n-1}([w^{(1)}_{k_1}F^{(1)}_{k_1}.w^{(2)}_1F^{(2)}_1w^{(2)}_2]),\ \sigma^{n-1}([w^{(2)}_2F^{(2)}_2.w^{(2)}_3F^{(2)}_3w^{(2)}_4]),...,$$

$$\sigma^{n-1}([w^{(2)}_{k_2-1}F^{(2)}_{k_2-1}.w^{(2)}_{k_2}F^{(2)}_{k_2}w^{(3)}_1]).$$

In particular, we obtain that $B(\Xi_n) \subseteq B(\Xi_{n-1})$ and $\{\Xi_n\}_{n\ge0}$ is a nested sequence of K-R partitions.

(3) We claim that the partitions $\{\Xi_n\}_{n\ge0}$ generate the topology of X_σ. Observe that it suffices to show that the function $x_{[-m,m]}$, $m > 0$, is constant on each element of partition $\{\Xi_n\}$ for n big enough. To see this, choose n such that

$$\min\{|\sigma^n(a)| \ : \ a \in A_l\} > m.$$

Fix a word $W = (v_1S_1v_2S_2v_3) \in \mathcal{V}$ and $0 \le k < |\sigma^n(v_2S_2)|$. For each $x \in T^k\sigma^n([v_1S_1.v_2S_2v_3])$, there exists $y \in [v_1S_1.v_2S_2v_3]$ such that $x = T^k\sigma^n(y)$. It follows that the word $\sigma^n(v_2S_2v_3)$ is a prefix of $\sigma^n(y[0,\infty))$ and $\sigma^n(v_1S_1)$ is a suffix of $\sigma^n(y(-\infty,-1])$. Therefore,

$$\sigma^n(y)[-L,R) = \sigma^n(v_1S_1v_2S_2v_3),$$

where $L = |\sigma^n(v_1S_1)|$ and $R = |\sigma^n(v_2S_2v_3)|$. Thus, we obtain

$$x_{[-m,m]} = \sigma^n(v_1S_1v_2S_2v_3)_{[L+k-m,L+k+m)},$$

which does not depend on x, but only on k and the word W.

Now it follows from Theorem 2.7 that the substitutional system (X_σ, T_σ) is homeomorphic to the Bratteli-Vershik system (X_B, φ_B), where the ordered Bratteli diagram B is constructed by the sequence of K-R partitions $\{\Xi_n\}_{n\geq 0}$. The fact that the diagram B is stationary has been proved in (2). □

Example 6.7. Consider the Chacon type substitution on the alphabet $A = \{0, s, 1\}$ given by $\sigma(0) = 00s0$, $\sigma(s) = s$, and $\sigma(1) = 0110$. Define the set $\mathcal{V}$ as in the proof of Theorem 6.6. Then $\mathcal{V}$ consists of the words $w_1, \ldots, w_8$, where

$$\begin{array}{llll} w_1 = 0.00 & w_2 = 0s.00 & w_3 = 0.0s0 & w_4 = 0s.0s0 \\ w_5 = 1.00 & w_6 = 0.11 & w_7 = 0.01 & w_8 = 1.10 \end{array}$$

Here the dot separates the negative and non-negative coordinates as in the definition of K-R partitions $\{\Xi_n\}$ in Theorem 6.6. To construct the Bratteli diagram, we need to trace the orbits of each base $\sigma^n[w_i]$, $i = 1, \ldots, 8$. In other words, we need to list all T_σ-towers which are intersected by $\sigma^n[w_i]$. This is achieved by decomposing each word $\sigma(w_i)$, $i = 1, \ldots, 8$, into the letters and analyzing the occurrence of long and short letters. We clarify the above scheme by considering the set $\sigma^n[w_4]$. Decompose $\sigma(w_4)$ into the letters:



This shows that the base $\sigma^n[w_4]$ consequently meets the bases $\sigma^{n-1}[w_2]$, $\sigma^{n-1}[w_3]$, and $\sigma^{n-1}[w_4]$. We denote this correspondence by $w_4 \mapsto w_2w_3w_4$. Repeating this argument for every set $\sigma^n([w_i])$, we get the following matching rule

$$\tau : \begin{cases} w_1 \mapsto w_1w_3w_2 \\ w_2 \mapsto w_2w_3w_2 \\ w_3 \mapsto w_1w_3w_4 \\ w_4 \mapsto w_2w_3w_4 \\ w_5 \mapsto w_1w_3w_2 \\ w_6 \mapsto w_7w_6w_8w_5 \\ w_7 \mapsto w_1w_3w_2 \\ w_8 \mapsto w_7w_6w_8w_5 \end{cases}$$

It follows from the proof of Theorem 6.6 that the system (X_σ, T_σ) is conjugate to the Vershik map of the following stationary ordered Bratteli diagram determined by the rule τ, i.e., τ is the substitution read from B.



Notice that by Theorem 6.4, the systems (X_σ, T_σ) and (X_τ, T_τ) are conjugate. See also Example 7.14 for another Bratteli-Vershik model of (X_σ, T_σ). We also mention the work [GjJo, Section 4.2] where the authors presented a Bratteli-Vershik model for the minimal component of the system (X_σ, T_σ).

7 Derivative substitutions

In the section, we study a generalization of the notion of derivative substitutions which was defined in [Dur] and [DHS] for primitive substitutions. We show that this notion works also for non-primitive substitutions.

To make the exposition clear and abandon some pathological situations, we restrict our study to the class of substitutions, which we call *m-primitive*. However, the results can be applied to more general class of substitutions, for example, for the Chacon type substitutions.

Definition 7.1. Let $m > 0$. We say that a substitution $\sigma : A \to A^+$ is *m-primitive* if we can decompose the alphabet $A = A_1 \sqcup \ldots \sqcup A_m \sqcup A_0$ with $|A_i| \geq 2$, $i = 1, \ldots, m$, and

(a) $\sigma(A_i) \subset A_i^+$ and $\sigma|_{A_i}$ is primitive for every $i = 1, \ldots, m$;

(b) for all $a \in A$ there exists a letter $b \in A_1 \cup \ldots \cup A_m$ and $i > 0$ with $b \prec \sigma^i(a)$.

(c) $L(\sigma) = L(\sigma^k)$ for every $k \geq 1$.

(d) $A \subset L(X_\sigma)$.

Given an m-primitive substitution $\sigma : A \to A^+$, define the matrix of the substitution $M(\sigma) = (M(\sigma)_{a,b})_{a,b\in A}$ as follows: $M(\sigma)_{a,b}$ is the number of occurrences of b in $\sigma(a)$. Then the matrix $M(\sigma)$ is of the form

$$M(\sigma) = \begin{pmatrix} M_1 & O & O & \dots & O & O & \dots & O \\ O & M_2 & O & \dots & O & O & \dots & O \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ O & O & O & \dots & M_m & O & \dots & O \\ p_{1,1} & p_{1,2} & p_{1,3} & \cdots & p_{1,m} & p_{1,m+1} & \cdots & p_{1,m+q} \\ p_{2,1} & p_{2,2} & p_{2,3} & \cdots & p_{2,m} & p_{2,m+1} & \cdots & p_{2,m+q} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ p_{n,1} & p_{n,2} & p_{n,3} & \cdots & p_{n,m} & p_{n,m+1} & \cdots & p_{n,m+q} \end{pmatrix} \tag{11}$$

where O denotes a zero matrix, the matrices M_i, $i = 1, \dots, m$, are primitive, and for each $j = 1, \dots, n$ there is a power k of M such that at least one of the entries $p^{(k)}_{j,1}, \dots, p^{(k)}_{j,m}$ from M^k is not zero. We notice also that $|\sigma^n| \to \infty$.

By Proposition 5.6 the number of minimal components of the substitutional system (X_σ, T_σ) is bounded by $|A|$. However, for the class of m-primitive substitutions, the set of all minimal components admits a complete description.

Proposition 7.2 *Let σ be an m-primitive substitution. Then (X_σ, T_σ) has exactly m minimal components, which are*

$$Z_i = \{x \in X_\sigma \ : \ x_{[-n,n]} \textit{ is a factor of some } \sigma^m(a),\ a \in A_i\},$$

$i = 1, \dots, m$.

Proof. The fact that Z_i, $i = 1, \dots, m$, are minimal components is well-known [Que]. We will show that there are no others. For each Z_i, we can find $w_i \in Z_i$ and $p_i > 0$ such that $\sigma^{p_i}(w_i) = w_i$, see for example [Que, Chapter V]. Without loss of generality, we may assume that $\sigma(w_i) = w_i$ for all $i = 1, \dots, m$. We also assume that all letters from A_i appear in $\sigma(a)$ for every $a \in A_i$, $i = 1, \dots, m$.

Take any $n > 0$. For every $x \in X_\sigma$, there is $a \in A$ such that the word $\sigma^{n+2}(a)$ appears in x. Find $b \in A_i$ that appears in $\sigma(a)$ for some $i = 1, \dots, m$. Therefore, the word $\sigma^n(w_i(0))$ appears in x. Find $1 \leq i_0 \leq m$ and an infinite set $I \subseteq \mathbb{N}$ such that the word $\sigma^n(w_{i_0}(0))$ appears in x for all $n \in I$. This implies that the closure of the orbit of x contains the point w_{i_0}. Hence, $\overline{Orb_{T_\sigma}(x)} \supseteq Z_{i_0}$. This proves the result. $\square$

Remark 7.3. Note that (X_σ, T_σ) has a periodic point iff (Z_i, T_σ) is periodic for some i. Observe also that there is an algorithm that decides whether σ has periodic points or not, see, for example, the book [Kur] or references in [DHS].

7.1 Return words and proper substitutions

Let $\sigma : A \to A^+$ be an m-primitive substitution. Here we assume that (X_σ, T_σ) has no periodic points. By Proposition 7.2, we get that the system (X_σ, T_σ) has exactly m minimal components $Z_1, \dots, Z_m$. By standard arguments, we can find $w_i \in Z_i$ and $p_i > 0$ such that $\sigma^{p_i}(w_i) = w_i$ [Que, Chapter V]. Without loss of generality, we can assume that $\sigma(w_i) = w_i$ for all $i = 1, \dots, m$. Notice also that $\overline{Orb_{T_\sigma}(w_i)} = Z_i$ for each $i = 1, \dots, m$.

Set $r_i = w_i[-1]$ and $l_i = w_i[0]$ for $i = 1, \dots, m$. Observe that each word $\sigma(r_i)$ ends with r_i, whereas every word $\sigma(l_i)$ begins with l_i, for all $i = 1, \dots, m$.

Consider the clopen sets $[r_i.l_i] := \{x \in X_\sigma : x[-1] = r_i,\ x[0] = l_i\}$ and define

$$W = \bigcup_{i=1}^{m} [r_i.l_i].$$

Proposition 7.4 *The set W meets each T_σ-orbit and consists of recurrent points. Furthermore, the return time to W is bounded.*

Proof. Take any $x \in X_\sigma$ and consider $Z(x) = \overline{Orb_{T_\sigma}(x)}$. Since $Z(x)$ is a T_σ-invariant closed subset of X_σ, it contains one of the minimal components Z_i, $i = 1, \dots, m$, say Z_{i_0}. Therefore, $[r_{i_0}.l_{i_0}]$ meets the orbit of x. Hence $X_\sigma = \bigcup_{n \in \mathbb{Z}} T_\sigma^n W$. By compactness of X_σ, we see that $X_\sigma = \bigcup_{n=0}^{k} T_\sigma^n W$ for some $k > 0$. This shows that W consists of recurrent points. $\square$

Definition 7.5. We say that $w \in L(X_\sigma)$ is a *return word* if there exists $1 \le i, j \le m$ such that

(i) $r_i w l_j \in L(X_\sigma)$;
(ii) the first and last letters of w are l_i and r_j, respectively;
(iii) no word from $\{r_1 l_1, \dots, r_m l_m\}$ appears in w.

Let $\mathcal{R}$ denote the set of all return words. In view of Proposition 7.4, the set $\mathcal{R}$ is finite. Let us enumerate the return words in an arbitrary way and denote $\mathcal{N} = \{1, 2, \dots, card(\mathcal{R})\}$. Let $\phi : \mathcal{N} \to \mathcal{R}$ be an "enumeration" map.

The proof of the following proposition is trivial, so we omit it.

Proposition 7.6 *The maps $\phi_1 : \mathcal{N}^+ \to A^+$ and $\phi_2 : \mathcal{N}^{\mathbb{Z}} \to A^{\mathbb{Z}}$ induced by ϕ are one-to-one.*

Take any return word $w \in \mathcal{R}$. Decompose it into letters $w = w_1 \dots w_k$. By definition of return words, $w_1 = l_i$ and $w_k = r_j$ for some i, j, and $r_i w l_j \in L(\sigma)$. Therefore, $\sigma(r_i)\sigma(w)\sigma(l_j) \in L(\sigma)$. Since $\sigma(r_i)$ ends with r_i, whereas every word $\sigma(l_i)$ begins with l_i, we have that $r_i \sigma(w) l_j \in L(\sigma)$. So, the word $\sigma(w)$ appears in the word $r_i \sigma(w) l_j$ between occurrences of $r_i.l_i$ and $r_j.l_j$. Thus, by Proposition 7.6, $\sigma(w) = d_1, \dots, d_q$ can be uniquely written as a concatenation of return words d_i.

Definition 7.7. Define the substitution $\tau : \mathcal{R} \to \mathcal{R}^+$ by $\tau(w) = d_1 \dots d_q$, where $d_1 \dots d_q$ is the unique decomposition of $\sigma(w)$ into return words. The substitution τ is called the *derivative substitution* of σ.

Notice that

$$\phi \circ \tau^m = \sigma^m \circ \phi.$$

The following result justifies the name of the derivative substitution.

Proposition 7.8 *The substitutional dynamical system* (X_τ, T_τ) *associated to* $\tau : \mathcal{R} \to \mathcal{R}^+$ *is homeomorphic to the system induced by* (X_σ, T_σ) *on the cylinder set* $W = \bigcup_{i=1}^m [r_i.l_i]$.

Proof. By Proposition 7.6, we have that $\phi : \mathcal{R}^{\mathbb{Z}} \to A^{\mathbb{Z}}$ is one-to-one.

(1) $\phi(X_\tau) \subset X_\sigma$. Indeed, consider any $x = (x_i) \in X_\tau$. By definition of X_τ, for every $n > 0$, there are $w \in \mathcal{R}$ and an integer $m > 0$ such that $x_{[-n,n]}$ is a factor of $\tau^m(w)$. It follows that $\phi(x_{[-n,n]})$ is a factor of $\phi \circ \tau^m(w) = \sigma^m \circ \phi(w)$. Since $\phi(w) \in L(\sigma)$, we get that $\phi(x_{[-n,n]}) \in L(\sigma)$ for every $n > 0$.

(2) It is clear that $\phi(X_\tau) \subseteq W$.

(3) $\phi(X_\tau) = W$. Indeed, take any $z \in W$. By Proposition 7.4, we can decompose z into a concatenation of return words $z = \ldots \phi(x[-1])|\phi(x[0]) \ldots$. We must show that $x = (x[i]) \in X_\tau$. By the definition of X_σ for any $n > 0$ there are $k > 0$ and $a \in A$ such that $\phi(x[-n,n]) \prec \sigma^k(a)$. Take a return word $w \in \mathcal{R}$ such that the letter a is a factor of $\phi(w)$. Therefore,

$$\phi(x[-n,n]) \prec \sigma^k(a) \prec \sigma^k(\phi(w)) = \phi(\tau^k(w)).$$

It follows from the definition of ϕ that $x[-n,n]$ is a factor of $\tau^k(w)$.

(4) If $x \in X_\tau$ and $z = \phi(x)$, then the first return time of z to W is $n = |\phi(x_0)|$. Thus, the image of z by the first return time transformation is $T_\sigma^n z = \phi(T_\tau x)$. This proves the proposition. □

Now we introduce the notion of a proper substitution. Our motivation to bring in this notion comes from the paper [DHS], where proper substitutions were indispensable for the Bratteli diagram construction. However, our definition of proper substitutions differs from that of [DHS]. The key concept which we put behind this notion is that for proper substitutions (for both definitions: ours and from [DHS]) the sequence of K-R partitions $\{\mathcal{P}_n\}$ defined in Corollary 5.18 generates the topology.

Definition 7.9. Let $\sigma : A \to A^+$ be a substitution. We say that σ is *proper* if there is $p > 0$ such that for every letter $a \in A$, the first letter of $\sigma^p(b)$, with $b \in A$ and $ab \in L(X_\sigma)$, does not depend on b; and the last letter of $\sigma^p(c)$, with $c \in A$ and $ca \in L(X_\sigma)$, does not depend on c. □

Remark 7.10. Observe that if a substitution is proper in the sense of [DHS], then it is also proper by our definition.

Proposition 7.11 *The derivative substitution* τ, *defined by an* m*-primitive substitution, is a proper aperiodic* m*-primitive substitution.*

Proof. Denote by $\mathcal{R}_i$ the set of all return words that appear in the fixed point w_i, $i = 1, \ldots, m$. We will identify the elements of $\mathcal{R}$ with their counterparts in $L(X_\sigma)$ (see Definition 7.5). Set also $\mathcal{R}_0 = \mathcal{R} \setminus (\mathcal{R}_1 \cup \ldots \cup \mathcal{R}_m)$. To verify that $\tau : \mathcal{R}_i \to \mathcal{R}_i^+$ is a primitive substitution, we refer the reader to the proof of Lemma 21 from [DHS].

Denote by v_i the return word that appears first (if we are counting rightwards) in the fixed point w_i, $i = 1, \ldots, m$. That is v_i is a prefix of $(w_i)_{[0,+\infty)}$. Find $n > 0$ such that

$$|\sigma^n| > \max_{i=1,\ldots,m} |v_i|. \tag{12}$$

It follows that $v_i l_i$ is a prefix of $\sigma^n(l_i)$. Take any return words w and w' with $ww' \in L(\tau)$. By definition, r_{i_0} is a suffix of w for some $i_0 = 1, \ldots, m$. Therefore, the word w' begins with l_{i_0}. So $\sigma^n(l_{i_0})$ is a prefix of $\sigma^n(w')$. This implies that $v_{i_0} l_{i_0}$ is a prefix of $\sigma^n(w')$. This means that v_{i_0} is the first return word in $\sigma^n(w')$. That is $\tau^n(w')$ begins with v_{i_0}. Thus, the first letter of $\tau^n(w')$ does not depend on w', but only on w.

The same argument works to show that the last letter of $\tau^n(w')$ with $w'w \in L(\tau)$ depends only on w for n large enough.

Now consider any $w \in \mathcal{R}_0$. Then the first letter of w is l_i for some $i = 1, \ldots, m$. It follows that $\tau^n(w)$ contains v_i, where n is as in (12). □

To prove the following result, we use Proposition 14 from [DHS].

Proposition 7.12 *Let $\sigma : A \to A^+$ be an aperiodic proper substitution such that $|\sigma^n| \to \infty$. Then the sequence of partitions $\{\mathcal{P}_n\}$ defined in Corollary 5.18 generates the topology of X_σ.*

Proof. Let $p > 0$ be an integer as in Definition 7.9 of proper substitutions. Given an integer $m > 0$, we claim that for n sufficiently large the function $x_{[-m,m]}$ is constant on each element of partition $\{\mathcal{P}_n\}$. Choose n so large that $|\sigma^{n-p}| > m$.

Fix $a \in A$ and $0 \le k < |\sigma^n(a)|$. For each $x \in T^k\sigma^n([a])$, there exists $y \in X_\sigma$ such that $y_0 = a$ and $x = T^k\sigma^n(y)$. The word $\sigma^n(a)\sigma^n(y_1)$ is a prefix of $\sigma^n(y_{[0,\infty)})$. By definition of proper substitutions, the first letter of $\sigma^p(y_1)$, say l, does not depend on y_1 (it depends only on the letter a). It follows that $\sigma^n(a)\sigma^{n-p}(l)$ is a prefix of $\sigma^n(y_{[0,\infty)})$. Similarly, there is $r \in A$ that depends only on the letter a (not on y) and such that $\sigma^{n-p}(r)$ is a suffix of $\sigma^n(y_{(-\infty,-1]})$. Therefore,

$$\sigma^n(y)_{[-R,L)} = \sigma^{n-p}(r)\sigma^n(a)\sigma^{n-p}(l),$$

where $R = |\sigma^{n-p}(r)|$ and $L = |\sigma^n(a)| + |\sigma^{n-p}(l)|$. Thus, we obtain

$$x_{[-m,m]} = \sigma^{n-p}(r)\sigma^n(a)\sigma^{n-p}(l)_{[R+k-m,R+k+m)},$$

which does not depend on x, but only on k and a. □

Now, we are ready to present a general approach for construction of Bratteli diagrams for m-primitive substitutions. To build a Bratteli-Vershik model for σ, it is sufficient to construct a Bratteli diagram for τ and then add some edges to the first level.

Theorem 7.13 *Let σ be an m-primitive aperiodic substitution over an alphabet A with derivative substitution τ. Let $B = (V, E, \le)$ be the stationary ordered*

Bratteli diagram built by the matrix of the substitution τ where τ is read on B. Suppose also that B has $|w|$ edges between the top vertex and the vertex defined by the return word $w \in \mathcal{R}$. Then B admits a continuous dynamics (X_B, T_B) which is homeomorphic to (X_σ, T_σ).

Proof. Let B' be the stationary Bratteli diagram built by the matrix of the substitution τ with simple edges between the top vertex and vertices of the first level. By Proposition 7.11 τ is a proper substitution. It follows from Proposition 7.12 and results of Section 2 that the Bratteli diagram B' admits continuous dynamics $(X_{B'}, \varphi_{B'})$ which is homeomorphic to (X_τ, T_τ). Then the application of Proposition 7.8 yields the result. $\square$

Example 7.14. Let σ be the Chacon type substitution defined in Example 6.7. Note that the system (X_σ, T_σ) has only one minimal component which is generated by the fixed point

$$w = \lim_n \sigma^n(0).\sigma^n(0) = \ldots 00s0s00s0.00s000s0s00s0\ldots$$

Though σ is not an m-primitive substitution, the technique developed in the section can still be applied to build a Bratteli-Vershik model of (X_σ, T_σ).

Consider the set of all return words $\mathcal{R}$. One can check that

$$\mathcal{R} = \{v_1 = 0,\ v_2 = 0s0,\ v_3 = 0s0s0,\ v_4 = 0110\}.$$

Find the unique decomposition of each $\sigma(v_i)$ into return words

$$\begin{aligned}
&\sigma(v_1) = 0|0s0 = v_1v_2\\
&\sigma(v_2) = 0|0s0s0|0s0 = v_1v_3v_2\\
&\sigma(v_3) = 0|0s0s0|0s0s0|0s0 = v_1v_3v_3v_2\\
&\sigma(v_4) = 0|0s0|0110|0110|0|0s0 = v_1v_2v_4v_4v_1v_2
\end{aligned}$$

Thus, the derivative substitution $\tau : \mathcal{R} \to \mathcal{R}^+$ is defined by $\tau(v_1) = v_1v_2$; $\tau(v_2) = v_1v_3v_2$; $\tau(v_3) = v_1v_3v_3v_2$; and $\tau(v_4) = v_1v_2v_4v_4v_1v_2$. Clearly, the substitution τ is proper and $|\tau^n| \to \infty$. By Theorem 7.13, the system (X_σ, T_σ) is conjugate to the Vershik map of the following stationary ordered Bratteli diagram:



Appendix: Description of the phase space X_σ

Here we give a combinatorial description of the phase space X_σ of a substitutional dynamical system assuming that $|\sigma^n(a)| \to \infty$ for all $a \in A$.

A. Denote by Λ the set of all sequences $\overline{s} = \{(a_n, b_n)\}_{n\geq 0}$ such that $a_n b_n \in L(\sigma)$ and a_n appears at the $(|\sigma(a_{n+1})| - 1)$-th position of $\sigma(a_{n+1})$ and b_n appears at zero position of $\sigma(b_{n+1})$.

B. Denote by $\mathcal{M}$ the set of all sequences $\overline{m} = \{(a_0, i_0), (a_1, i_1), \ldots\}$ such that $a_j \in A$ and i_j is a place of occurrence of a_{j-1} in $\sigma(a_j)$, $i_j \in \{0, 1, \ldots, |\sigma^n(a_j)| - 1\}$. We assume that $i_0 = 0$.

For a fixed $\overline{m}$, define inductively a sequence $\{j_n\}$ as follows: $j_0 = i_0 = 0$ and

$$j_{n+1} = \left\{ \begin{array}{ll} |\sigma^n(a_0^{(n+1)})| + \ldots + |\sigma^n(a_{i_{n+1}-1}^{(n+1)})| + j_n & \text{if } i_{n+1} \geq 1, \\ j_n & \text{if } i_{n+1} = 0, \end{array} \right. \tag{A.1}$$

here $\sigma(a_{n+1}) = a_0^{(n+1)} a_1^{(n+1)} \ldots a_{|\sigma(a_{n+1})|-1}^{(n+1)}$ with $a_i^{(n+1)} \in A$.

C. Denote by $\mathcal{M}_0$ the set of all $\overline{m}$ for which $j_n \to \infty$ and $(|\sigma^n(a_n)| - j_n) \to \infty$ as $n \to \infty$.

Construction of sequences from X_σ

- For every sequence $\overline{s} = \{(a_n, b_n)\}_{\geq 0} \in \Lambda$, define the sequence $w = w(\overline{s}) \in X_\sigma$ by

$$w_{[-|\sigma^n(a_n)|, |\sigma^n(b_n)|-1]} = \sigma^n(a_n).\sigma^n(b_n) \text{ for all } n \geq 0. \tag{A.2}$$

- For every $\overline{m} \in \mathcal{M}_0$, define the sequence $w = w(\overline{m})$ by

$$w_{[-j_n, |\sigma^n(a_n)|-j_n-1]} = \sigma^n(a_n) \tag{A.3}$$

(it is easily seen that the sequences $w(\overline{s})$ and $w(\overline{m})$ indeed belong to X_σ).

Let us recall that $T_\sigma : A^{\mathbb{Z}} \to A^{\mathbb{Z}}$ denote the shift. The next theorem describes the phase space of an arbitrary substitution σ.

Theorem A.1 *Let $\sigma : A \to A^+$ be a substitution with $|\sigma^n| \to \infty$. (1)*

$$X_\sigma = \{w(\overline{m}) : \overline{m} \in \mathcal{M}_0\} \cup \bigcup_{\overline{s} \in \Lambda} Orb_{T_\sigma}(w(\overline{s})) \tag{A.4}$$

(2) the set $\{\omega(\overline{s}) : \overline{s} \in \Lambda\}$ is finite.
(3) If σ is aperiodic, then

$$\bigcap_{n=0}^{\infty} \sigma^n(X_\sigma) = \{w(\overline{s}) \ : \ \overline{s} \in \Lambda\}.$$

Proof. (1) It follows from Remark 5.11 that for every $t \in X_\sigma$, there exists a sequence $\mathcal{F} = \{(x_n, (y_n, i_n), z_n)\}_{n \geq 0}$ such that

i. $x_n, y_n, z_n \in A$ and $x_n y_n z_n \in L(\sigma)$;

ii. $i_n \in \{0, \ldots, |\sigma^n(y_n)| - 1\}$ is the position of occurrence of y_{n-1} within $\sigma(y_n)$;

iii. If $i_n = 0$, then x_{n-1} appears in $\sigma(x_n)$ at the position $|\sigma(x_n)| - 1$;

iv. If $i_n = |\sigma(y_n)| - 1$, then z_{n-1} appears in $\sigma(z_n)$ at zero position;

v. If $0 < i_n < |\sigma(y_n)| - 1$, then $x_{n-1} y_{n-1} z_{n-1}$ appears in $\sigma(y_n)$ at the position $i_n - 1$.

vi. If the sequence $\{j_n\}$ is determined by the sequence $\overline{m} = \{(y_n, i_n)\}_{n \geq 0} \in \mathcal{M}$ as in (A.1), then

$$t[-j_n - |\sigma^n(x_n)|, |\sigma^n(y_n)| - j_n - 1 + |\sigma^n(z_n)|] = \sigma^n(x_n)\sigma^n(y_n)\sigma^n(z_n). \tag{A.5}$$

Now we have three options:

(a) $j_n \to \infty$ and $|\sigma^n(y_n)| - j_n \to \infty$ as $n \to \infty$. In this case, we get that $t = w(\overline{m})$.

(b) The sequence $\{j_n\}$ is bounded and $(|\sigma^n(y_n)| - j_n) \to \infty$. Here, we get that $w = w(\overline{m})$ is one-sided (to the right). Since the sequence $\{j_n\}$ is non-decreasing and bounded, there is $n_0 > 0$ such that $j_n = j_{n_0}$ for all $n \geq n_0$. This implies that $i_n = 0$ for all $n \geq n_0$, i.e. y_n appears at zero position in $\sigma(y_{n+1})$. This, in particular, implies that x_n appears in $\sigma(x_{n+1})$ at the last position. Therefore,

$$t[-j_{n_0} - |\sigma^n(x_n)|, |\sigma^n(y_n)| - j_{n_0} - 1] = \sigma^n(x_n)\sigma^n(y_n) \text{ for } n \geq n_0.$$

Set $x'_{n_0-1} = \sigma(x_{n_0})[|\sigma(x_{n_0})| - 1]$ and $y'_{n_0-1} = \sigma(y_{n_0})[0]$. Define inductively $x'_{k-1} = \sigma(x'_k)[|\sigma(x'_k)| - 1]$ and $y'_{k-1} = \sigma(y'_k)[0]$ for $k = n_0 - 1, \ldots, 1$. Setting

$$\overline{s} = \{(x'_0, y'_0), \ldots, (x'_{n_0-1}, y'_{n_0-1}), (x_{n_0}, y_{n_0}), (x_{n_0+1}, y_{n_0+1}), \ldots\},$$

we get that $t = T^{j_0}(w(\overline{s}))$.

(c) The sequence $\{|\sigma^n(y_n)| - j_n\}$ is bounded and $j_n \to \infty$ as $n \to \infty$. The proof in this case is similar to (b).

(2) Denote by A_0 (by A_1) the set of all letters $a \in A$ such that $\sigma^{n_a}(a)$ begins (ends) with a for some n_a. Clearly, if such an n_a exists, then it can be chosen from the interval $[1, |A| + 1]$. Set $p = \prod_{a \in A_0} n_a \times \prod_{b \in A_1} n_b$. Then p is bounded by $(|A| + 1)^{2|A|}$.

Consider a sequence $\overline{s} = \{(a_n, b_n)\}_{n \geq 0} \in \Lambda$. Find an infinite set I such that $a_k = a_{k'}$ and $b_k = b_{k'}$ for all $k, k' \in I$. Therefore, $\sigma^{k-k'}(a_k)$ ends with a_k for all $k > k'$, $k, k' \in I$. This shows that $\sigma^p(a_k)$ $(\sigma^p(b_k))$ ends (begins) with a_k (b_k) for all $k \in I$. We can write down each $k \in I$ as

$$k = pl_k + m_k \text{ with } l_k \geq 0 \text{ and } 0 \leq m_k < p.$$

Find an infinite set $J \subseteq I$ such that $m_k = m_{k'}$ for $k, k' \in J$. It follows that $a_{m_k + pn} = a_{m_k}$ and $b_{m_k + pn} = b_{m_k}$ for all $n \geq 0$ and $k \in J$. Then

$$w(\overline{s}) = \lim_{n \to \infty} \sigma^n(a_n).\sigma^n(b_n) = \lim_{n \to \infty} \sigma^{pn}(\sigma^{m_k}(a_{m_k})).\sigma^{pn}(\sigma^{m_k}(b_{m_k})),$$

where k is any integer from J. This shows that each element $w(\overline{s})$ is determined by a finite number of parameters taken from finite sets.

(3) If $y \in Y = \bigcap_{n \geq 0} \sigma^n(X_\sigma)$, then for every $n \geq 0$ we have $y = \sigma^n(x_n)$ for some $x_n \in X_\sigma$. By Theorem 5.17, the point x_n is uniquely defined. Therefore, $\sigma(x_{n+1}) = x_n$ for every n. Setting $\overline{s} = \{(x_n[-1], x_n[0])\}_{n \geq 0}$, we get that $y = w(\overline{s})$.

Conversely, if $y = w(\overline{s})$, then it is not hard to decompose y into n-words such that $y = \sigma^n(x_n)$ for some $x_n \in X_\sigma$. In particular, this shows that $y \in Y$.□

Acknowledgments. The work was done during our mutual visits to the University of Torun and Institute for Low Temperature Physics. We are thankful to these institutions for the hospitality and support. We would like to express our special thanks to C. Skau and B. Solomyak for numerous stimulating discussions and to T. Downarowicz and A. Maass for sending us their paper and drawing our attention to their result. K. Medynets also acknowledges the support of INTAS Young Scientist Fellowship.

References

[BDM] S. Bezuglyi, A. H. Dooley, and K. Medynets. The Rokhlin lemma for homeomorphisms of a Cantor set. *Proc. Am. Math. Soc.*, 133(10):2957 – 2964, 2005.

[CE1] T. Carlsen and S. Eilers. Augmenting dimension group invariants for substitution dynamics. *Ergod. Th. & Dynam. Sys.*, 24(4):1015–1039, 2004.

[CE2] T. Carlsen and S. Eilers. Ordered k-groups associated to substitutional dynamics. *J. Funct. Anal.*, 238(1):99–117, 2006.

[DHS] F. Durand, B. Host, and B. Skau. Substitutional dynamical systems, Bratteli diagrams and dimension groups. *Ergod. Th. & Dynam. Sys.*, 19(4):953–993, 1999.

[DM] T. Downarowicz and A. Maass. Finite rank Bratteli-Vershik diagrams are expansive. to appear in *Ergod. Th. & Dynam. Sys.*.

[Dur] F. Durand. A characterization of substitutive sequences using return words. *Discrete Math.*, 179:89–101, 1998.

[Fer] S. Ferenczi. Substitution dynamical systems on infinite alphabets. *Preprint*, 2005.

[Fog] Pytheas N. Fogg. *Substitutions in dynamics, arithmetics and combinatorics*, volume 1794 of *Lecture Notes in Mathematics*. Springer, Berlin, 2002.

[For] A.H. Forrest. K-groups associated with substitution minimal systems. *Israel J. Math.*, 98:101 – 139, 1997.

[GjJo] R. Gjerde and Ø. Johansen. Bratteli-Vershik models for Cantor minimal systems: applications to Toeplitz flows. *Ergod. Th. & Dynam. Sys.*, 20:1687–1710, 2000.

[Gla] E. Glasner. *Ergodic Theory via Joinings*, volume 101 of *Mathematical Surveys and Monographs*. AMS, 2003.

[GMPS] T. Giordano, H. Matui, I. Putnam, and C. Skau. Orbit equivalence for cantor minimal Z^2-systems. *Preprint*, 2006.

[GPS1] T. Giordano, I. Putnam, and C. Skau. Topological orbit equivalence and C^*-crossed products. *J. Reine Angew. Math.*, 469:51 – 111, 1995.

[GPS2] T. Giordano, I. Putnam, and C. Skau. Full groups of Cantor minimal systems. *Israel. J. Math.*, 111:285–320, 1999.

[GPS3] T. Giordano, I. Putnam, and C. Skau. Affable equivalence relations and orbit structure of Cantor dynamical systems. *Ergodic Theory Dyn. Syst.*, 24(2):441–475, 2004.

[GW] E. Glasner and B. Weiss. Weak orbit equivalence of Cantor minimal systems. *Int. J. Math.*, 6(4):569–579, 1995.

[Hos] B. Host. Valeurs propres des systémes dynamiques dédinis par des substitutions de longueur variable. *Ergod. Th. & Dynam. Sys.*, 6(4):529–540, 1986.

[HPS] R. H. Herman, I. Putnam, and C. Skau. Ordered bratteli diagrams, dimension groups, and topological dynamics. *Int. J. Math.*, 3(6):827 – 864, 1992.

[Kur] P. Kurka. *Topological and Symbolic Dynamics.* Cours Spécialisés (Paris) 11. Société Mathématique de France, Paris, 2003.

[Med] K. Medynets. Cantor aperiodic systems and Bratteli diagrams. *C. R., Math., Acad. Sci. Paris*, 342(1):43–46, 2006.

[Mos1] B. Mossé. Puissances de mots et reconnaissabilite des points fixes d'une substitution. *Theor. Comput. Sci.*, 99(2):327–334, 1992.

[Mos2] B. Mossé. Reconnaissabilite des substitutions et complexite des suites automatiques. *Bull. Soc. Math. Fr.*, 124(2):329–346, 1996.

[Que] M. Queffelec. *Substitution dynamical systens — spectral analysis*, volume 1294 of *Lect. Notes in Math.* Springer-Verlag, 1987.

[Wal] P. Walters. *An Introduction to Ergodic Theory*, volume 79 of *Graduate Texts in Mathematics*. Springer, Berlin, 1982.

[Yua1] H. Yuasa. On the topological orbit equivalence in a class of substitutional minimal systems. *Tokyo, J. Math.*, 25(2):221–240, 2002.

[Yua2] H. Yuasa. Invariant measures for the subshifts arising from non-primitive substitutions. *Preprint*, 2006.